

Human Monocyte Chemoattractant Protein-1(rDNA derived) Reference standard

Cat. No. CCL2-125H Lot. No. (See product label)

SPECIFICATION

Product Overview

Species Human

Enzyme Unit The assigned potency is 5000 arbitrary units per ampoule.

Usage

No attempt should be made to weigh out any portion of the freeze-dried material prior to reconstitution. Dissolve the total contents of the ampoule in 0.5ml of sterile distilled water. Rinse the ampoule with a further 0.4ml of sterile distilled water and make up the total volume to 1.0ml. This solution will contain MCP-1 at a concentration of 5,000 units per ml. It is advisable to include a carrier protein during initial reconstitution to prevent loss of material if this does not interfere in the intended assay method.

Notes

This preparation is not for administration to humans. The material is not of human or bovine origin. As with all materials of biological origin, this preparation should be regarded as potentially hazardous to health. It should be used and discarded according to your own laboratory's safety procedures. Such safety procedures should include the wearing of protective gloves and avoiding the generation of aerosols. Care should be exercised in opening ampoules or vials, to avoid cuts.

Stability

Reference materials are held at Creative Biomart within assured, temperature-controlled storage facilities and they should be stored on receipt as indicated on the label. An arbitrary unitage was assigned to this preparation at the time of production without confirmation of long term stability. Evidence from similar materials prepared

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

by an equivalent process indicates that long term stability is likely to be maintained and the material is suitably stable for shipment at ambient temperature without any effect on the assigned values. Once reconstituted, diluted or aliquoted, users should determine the stability of the material according to their own method of preparation, storage and use. Users who have data supporting any deterioration in the characteristics of any reference preparation are encouraged to contact Creative Biomart.

Storage

For economy of use, it is recommended that the solution be sub divided into aliquots and stored at +4°C for up to 1 week. Unopened ampoules should be stored at –20°C. Please note: because of the inherent stability of lyophilized material, Creative Biomart may ship these materials at ambient temperature.

GENE INFORMATION

Gene Name CCL2 chemokine (C-C motif) ligand 2 [Homo sapiens];

Official Symbol CCL2

Synonyms

CCL2; chemokine (C-C motif) ligand 2; SCYA2, small inducible cytokine A2 (monocyte chemotactic protein 1, homologous to mouse Sig je); C-C motif chemokine 2; GDCF 2; HC11; MCAF; MCP 1; MCP1; MGC9434; monocyte chemoattractant protein 1; monocyte chemotactic and activating factor; monocyte chemotactic protein 1; homologous to mouse Sig je; monocyte secretory protein JE; small inducible cytokine subfamily A (Cys Cys); member 2; SMC CF; small-inducible cytokine A2; monocyte chemoattractant protein-1; small inducible cytokine subfamily A (Cys-Cys), member 2; small inducible cytokine A2 (monocyte chemotactic protein 1, homologous to mouse Sig-je); MCP-1; SCYA2; GDCF-2; SMC-CF; HSMCR30;

Gene ID 6347

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA Refseq	NM_002982
Protein Refseq	NP_002973
UniProt ID	P13500
Chromosome Location	17q11.2-q21.1
Pathway	Activation of Genes by ATF4, organism-specific biosystem; Chagas disease (American trypanosomiasis), organism-specific biosystem; Chagas disease (American trypanosomiasis), conserved biosystem; Chemokine receptors bind chemokines, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Class A/1 (Rhodopsin-like receptors), organism-specific biosystem;
Function	CCR2 chemokine receptor binding; CCR2 chemokine receptor binding; G-protein coupled receptor binding; chemokine activity; heparin binding; protein kinase activity; receptor binding;