

Recombinant Human PSMD4, GST-tagged

Cat. No. PSMD4-152H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human PSMD4 (amino acid residues 1-377), with N-terminal GST, was expressed in E.coli.
Species	Human
Source	E.coli
ProteinLength	1-377 a.a.
Description	S5a (Rpn10) is a major ubiquitin binding protein that binds preferentially to polyubiquitin chains. It is found as a subunit of the 26S proteasome, but unlike other proteasome subunits, S5a also exists predominantly as a free protein in the cytosol. S5a contains two stretches of approximately 15 amino acids called the ubiquitin interacting motifs (UIMs), which are responsible for its high affinity for ubiquitin chains (Kim and Goldberg, 2012).
Form	50 mM HEPES pH 7.5, 150 mM sodium chloride, 2 mM dithiothreitol, 10% glycerol
Molecular Mass	~67.6kDa
Purity	>98% by InstantBlue™ SDS-PAGE
Storage	12 months at -70°C. Avoid multiple freeze/thaw cycles.
Concentration	0.5 mg/ml

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

GENE INFORMATION

Gene Name	PSMD4 proteasome (prosome, macropain) 26S subunit, non-ATPase, 4 [Homo sapiens]
Official Symbol	PSMD4
Synonyms	PSMD4; proteasome (prosome, macropain) 26S subunit, non-ATPase, 4; 26S proteasome non-ATPase regulatory subunit 4; AF; AF 1; Rpn10; S5A; angiocidin; RPN10 homolog; antisecretory factor 1; S5a/antisecretory factor protein; multiubiquitin chain-binding protein; 26S proteasome regulatory subunit S5A; ASF; AF-1; MCB1; pUB-R5;
Gene ID	5710
mRNA Refseq	NM_002810
Protein Refseq	NP_002801
MIM	601648
UniProt ID	P55036
Chromosome Location	1q21.2
Pathway	APC/C-mediated degradation of cell cycle proteins, organism-specific biosystem; APC/C:Cdc20 mediated degradation of Securin, organism-specific biosystem; APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem; APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cdh1 targeted proteins in late mitosis/early G1, organism-specific biosystem; Activation of APC/C

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

and APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem; Activation of NF-kappaB in B Cells, organism-specific biosystem; Adaptive Immune System, organism-specific biosystem;

Function

protein binding;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA