

Recombinant Mouse Tnf protein

Cat. No. Tnf-484M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Mouse Tnf protein was expressed in Escherichia coli.
Species	Mouse
Source	E.coli
ProteinLength	157
Description	Tumor necrosis factor alpha (TNF-α), also called cachectin, is the best-know member of the TNF-family, which can cause cell death. This protein is produced by neutrophils, activated lymphocytes, macrophages, NK cells, LAK cells, astrocytes endothelial cells, smooth muscle cells and some transformed cells. TNF-α occurs as a secreted, soluble form and as a membrane-anchored form, both of which are biologically active. The naturally-occurring form of TNF-α is glycosylated, but non-glycosylated recombinant TNF-α has comparable biological activity. The biologically active native form of TNF-α is reportedly a trimer. Human and murine TNF-α show approximately 79 % homology at the amino acid level and cross-reactivity between the two species. Two types of receptors for TNF-α have been described and virtually all cell types studied show the presence of one or both of these receptor types.
Form	Lyophilized from a 0.2 μ m filtered solution in PBS, pH 7.2.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by a cytotoxicity assay using murine L929 cells is less than 0.1 ng/ml, corresponding to a

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	specific activity of $> 1.0 \times 10^7$ IU/mg in the presence of actinomycin D.
Molecular Mass	Approximately 17.4 kDa. The recombinant murine TNF- α is a soluble 157 amino acid protein which corresponds to C-terminal extracellular domain of the full length transmembrane protein.
AA Sequence	MLRSSSQNSSDKPVAHVVAHQVEEQLEWLSQRANALLANGMDLKDNQLVVPADG LYLVYSQVLFGKGQCPDYVLLTHTVSRFAISYQEKVNLLSAVKSPCPKDTPEGAELK PWYEPIYLGGVFQLEKGDQLSAEVNLPKYLDFAESGQVYFGVIAL
Endotoxin	Less than 1 EU/ μ g of rMuTNF- α as determined by LAL method.
Purity	>98% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	Tnf
Official Symbol	Tnf

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Synonyms	TNF; tumor necrosis factor; TNF-a; TNF alpha; cachectin; tumor necrosis factor alpha; tumor necrosis factor-alpha; tumor necrosis factor ligand superfamily member 2; DIF; Tnfa; TNFSF2; Tnfsf1a; TNFalpha; TNF-alpha; MGC151434;
Gene ID	21926
mRNA Refseq	NM_013693
Protein Refseq	NP_038721
UniProt ID	P06804
SDS-PAGE of Tnf-484M	