

Human Foreskin Fibroblast Whole Cell Lysate

Cat. No. Foreskin Fibroblast-5H **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species Human

Concentration 2 mg/ml

Tissue Type Human Foreskin Fibroblast

Preparation method

The cells were grown in DMEM supplemented with 10% FBS (Fetal Bovine Serum). The lysate was prepared by first washing the cells in PBS. Washed cells were then incubated on ice in modified RIPA buffer containing 150 mM sodium chloride, 50 mM Tris HCl, pH 7.4, 1 mM EDTA, 1.0% NP-40, 0.5% sodium deoxycholic acid and 0.1% SDS to lyse the cells. Protein integrity was ensured using a cocktail of protease inhibitors with broad specificity for the inhibition of aspartic, cysteine, and serine proteases as well as aminopeptidases (0.1 mM AEBSF HCl, 0.08 M Aprotinin, 5 M Bestatin, 1.5 M E-64, 2 M Leupeptin Hemisulfate, 1 M Pepstatin A). Phosphatase inhibitors 1 mM NaF and 1 mM Na₃VO₄ were also added. Cell debris was removed by centrifugation. Protein concentration was determined by modified Lowry assay using a commercially available kit. Protein concentration was adjusted to 2 mg/ml and then an equal volume of 2X SDS-PAGE sample buffer was added.

Recommended Usage

For research use only, not for diagnostic or therapeutic use.

Storage Buffer

1X SDS-PAGE Sample Buffer (62.5 mM Tris HCl, 2% SDS, 10% Glycerol and

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

0.005% bromophenol blue, pH 6.8)

Applications

Protein Lysate for WB. Ready-to-use lysates are especially prepared as positive controls for separation by SDS-PAGE and subsequent western blot analysis. Lysates are prepared in denaturing buffer WITHOUT dissociating agents (i.e. no 2-mercaptoethanol or dithiothreitol has been added).

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA