

Recombinant Human Interleukin 17C, Fc Chimera

Cat. No. IL17C-280H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human Interleukin 17C is produced in mammalian cells and fused with human IgG1Fc. IL-17C (C-terminus fused with human IgG1Fc) contains 408 amino acids with predicted MW 45.43 kDa x 2 (dimer) and 14 cysteines (Fc has 6) with no N-link glycosylation site. It stimulates the release of tumor necrosis factor alpha and IL-1beta from the monocytic cell line THP-1.
Species	Human
Source	Mammalian Cells
Description	Interleukin-17 (IL-17, or IL-17A) is the founding member of a group of cytokines called the IL-17 family. IL-17A, was originally identified as a transcript from a rodent T-cell hybridoma by Rouvier et al. in 1993. Known as CTLA8 in rodents, IL-17 shows high homology to viral IL-17 encoded by an open reading frame of the T lymphotropic rhadinovirus Herpesvirus saimiri. To elicit its functions, IL-17 binds to a type I cell surface receptor called IL-17R of which there are at least three variants IL17RA, IL17RB, and IL17RC.
Purity	>90%.
Quantity	2.0mg/2ml PBS.
Availability	Back order, one month for production.
Storage	-70°C.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

GENE INFORMATION

Gene Name	IL17C interleukin 17C [Homo sapiens]
Synonyms	IL17C; interleukin 17C; CX2; IL-21; IL-17C; MGC126884; MGC138401; interleukin-17C; cytokine CX2
Gene ID	27189
mRNA Refseq	NM_013278
Protein Refseq	NP_037410
MIM	604628
UniProt ID	Q9P0M4
Chromosome Location	16q24
Function	cytokine activity

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA