

Active Recombinant Swine Tumor Necrosis Factor (TNF Superfamily, Member 2)

Cat. No. TNF-369S **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Swine TNF- α was expressed in Yeast. MW= 16.9 kDa (154 aa).
Species	Pig
Source	Yeast
Description	TNF- α is a homotrimer with a subunit molecular mass of 17 kDa and that it plays a major role in growth regulation, differentiation, inflammation, viral replication, tumorigenesis, and autoimmune diseases; and in viral, bacterial, fungal, and parasitic infections. Besides inducing hemorrhagic necrosis of tumors, TNF was found to be involved in tumorigenesis, tumor metastasis, viral replication, septic shock, fever, inflammation, and autoimmune diseases including Crohn's disease, and rheumatoid arthritis as well as graft-versus-host disease.
Formulation/ Reconstitution	Lyophilized. Reconstitute with sterile phosphate-buffered saline containing at least 0.1% carrier protein.
Purification	Ion-exchange chromatography.
Purity	>95% as visualized by SDS-PAGE analysis.
Amino Acid Sequence	SSSQTSDDKPV AHVVANVKAE GQLQWQSGYA NALLANGVKL KDNQLVVPTD GLYLIYSQVL FRGQGCPSTN VFLTHTISR AVSYQTKVNL LSAIKSPCQR ETPEGAEAKP WYEPIYLGGV FQLEKDDRLS AEINLPDYLD FAESGQVYFG IIAL.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Bioactivity The biological activity of recombinant swine TNF alpha was measured in a cytotoxicity assay using L929 mouse fibrosarcoma cells. The ED50 for this effect is typically 1 3 ng/mL.

Stability And Storage Stable for up to twelve months from date of receipt at -20°C. Stable for at least 3 months when stored in working aliquots with a carrier protein at -20°C. Avoid repeated freeze/thaw cycles.

GENE INFORMATION

Gene Name TNF tumor necrosis factor (TNF superfamily, member 2) [*Sus scrofa*]

Synonyms tumor necrosis factor (TNF superfamily, member 2); TNF; TNFa; tumor necrosis factor; TNF-a; TNF-alpha; cachectin; tumor-necrosis factor; tumor necrosis factor alpha; tumor necrosis factor ligand superfamily member 2

Gene ID 397086

mRNA Refseq NM_214022.1

Protein Refseq NP_999187.1

Chromosome Location 7p11-q11

UniProt ID P23563

Pathway Adipocytokine signaling pathway; Allograft rejection; Alzheimer's disease; Amyotrophic lateral sclerosis (ALS); Apoptosis; Asthma; Cytokine-cytokine receptor interaction; Dilated cardiomyopathy; Fc epsilon RI signaling pathway; Graft-versus-host disease; Hematopoietic cell lineage; Hypertrophic cardiomyopathy (HCM); MAPK

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

signaling pathway; NOD-like receptor signaling pathway; Natural killer cell mediated cytotoxicity; RIG-I-like receptor signaling pathway; Systemic lupus erythematosus; T cell receptor signaling pathway; TGF-beta signaling pathway; Toll-like receptor signaling pathway; Type I diabetes mellitus; Type II diabetes mellitus

PDB rendering based
on 1TNF.

