

Recombinant Human Chemokine (C-C Motif)Ligand 7

Cat. No. CCL7-36H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human Monocyte Chemotactic Protein-3 produced in E.Coli is a non-glycosylated, Polypeptide chain containing 76 amino acids and having a molecular mass of 9011 Dalton. The MCP-3 is purified by proprietary chromatographic techniques.
Species	Human
Source	E.coli
Description	Chemokine (C-C motif) ligand 7 (CCL7) is a small cytokine known as a chemokine that was previously called monocyte-specific chemokine 3 (MCP3). Due to CCL7 possessing two adjacent N-terminal cysteine residues in its mature protein, it is classified among the subfamily of chemokines known as CC chemokines. CCL7 specifically attracts monocytes, and regulates macrophage function. It is produced by certain tumor cell lines and by macrophages. This chemokine is located on chromosome 17 in humans, in a large cluster containing many other CC chemokines and is most closely related to CCL2 (previously called MCP1).
Amino Acid Sequence	The sequence of the first five N-terminal amino acids was determined and was found to be Gln-Pro-Val-Gly-Ile.
Physical Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Purity	Greater than 98.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Formulation	The protein was lyophilized from a concentrated (1mg/ml) sterile solution containing no additives.
Solubility	It is recommended to reconstitute the lyophilized Monocyte Chemotactic Protein-3 in sterile 18MΩ-cm H ₂ O not less than 100g/ml, which can then be further diluted to other aqueous solutions.
Biological Activity	The specific activity as determined by the ability of MCP-3 to chemoattract human peripheral blood at 10-100ng/ml.
Storage	Lyophilized MCP-3 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CCL7 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

GENE INFORMATION

Gene Name	CCL7 chemokine (C-C motif) ligand 7 [Homo sapiens]
Synonyms	CCL7; chemokine (C-C motif) ligand 7; FIC; MARC; MCP3; NC28; MCP-3; SCYA6; SCYA7; MGC138463; MGC138465; monocyte chemotactic protein 3; monocyte chemoattractant protein 3; small inducible cytokine A7
Gene ID	6354
mRNA Refseq	NM_006273
Protein Refseq	NP_006264
MIM	158106

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	P80098
Chromosome Location	17q11.2-q12
Pathway	Chemokine signaling pathway; Cytokine-cytokine receptor interaction
Function	chemokine activity; heparin binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA