

Recombinant Human HDAC2 Protein, His tagged

Cat. No. HDAC2-3565H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human HDAC2 protein was expressed with c-terminal His-tag in high-5 cells using baculovirus expression system and purified by using conventional chromatography techniques.
Species	Human
Source	Insect Cells
ProteinLength	1-488aa
Description	This gene product belongs to the histone deacetylase family. Histone deacetylases act via the formation of large multiprotein complexes, and are responsible for the deacetylation of lysine residues at the N-terminal regions of core histones (H2A, H2B, H3 and H4). This protein forms transcriptional repressor complexes by associating with many different proteins, including YY1, a mammalian zinc-finger transcription factor. Thus, it plays an important role in transcriptional regulation, cell cycle progression and developmental events. Alternative splicing results in multiple transcript variants.
Form	Liquid in. 20mM Tris-HCl buffer (pH 8.0) containing 20% glycerol, 0.1M NaCl, 1mM DTT, 0.1mM PMSF
Molecular Mass	56.4 kDa (496aa)
AASequence	MAYSQGGGKKKVCYYYDGDIGNYYYGQGHMPKPHRIRMTHNLLNLYGLYRKMEIY

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

RPHKATAEEMTKYHSDEYIKFLRSIRPDNMSEYSKQMQRFNVGEDCPVFDGLFEFC
 QLSTGGSVAGAVKLNRRQQTDMAVNWAGGLHHAKKSEASGFCYVNDIVLAILELLKY
 HQRVLYIDIDIHHGDGVVEAFYTTDRVMTVSFHKYGEYFPGTGDLRDIGAGKGKYYA
 VNFPMRDGIDDESYGQIFKPIISKVMEMYQPSAVVLQCGADSLSGDRLGCFNLTVKG
 HAKCVEVVKTFNLPLLMLGGGGYTIRNVARCWTYETAVALDCEIPNELPYNDYFEYF
 GPDKLHISPSNMTNQNTPEYMEKIKQRLFENLRMLPHAPGVQMQAIPEDAVHEDS
 GDEDGEDPDKRISIRASDKRIACDEEFSDSEDEGEGERNRVADHKKGAKKARIEED
 KKETEDKKTVDKKEEDKSKDNSGEKTDTKGKSEQLSNP

Endotoxin	< 1 EU/μg of protein (determined by LAL method)
Purity	> 85% by SDS-PAGE
Applications	SDS-PAGE
Notes	For research use only. This product is not intended or approved for human, diagnostics or veterinary use.
Storage	Can be stored at +2 to +8 centigrade for 1 week. For long term storage, aliquot and store at -20 to -80 centigrade. Avoid repeated freezing and thawing cycles.
Concentration	0.25 mg/mL (determined by Bradford assay)

GENE INFORMATION


Gene Name	HDAC2 histone deacetylase 2 [Homo sapiens (human)]
Official Symbol	HDAC2
Synonyms	HDAC2; histone deacetylase 2; RPD3; YAF1; YY1-associated factor 1; transcriptional regulator homolog RPD3; HD2

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID	3066
mRNA Refseq	NM_001527
Protein Refseq	NP_001518
MIM	605164
UniProt ID	Q92769

SDS-PAGE


3 µg by SDS-PAGE under reducing condition and visualized by coomassie blue stain.