

Recombinant Human Prolactin Receptor, GST-tagged

Cat. No. PRLR-58H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human Prolactin Receptor produced in <i>E. Coli</i> is a non-glycosylated, Polypeptide chain amino acids 432-623 and having a molecular mass of 45 kDa, the PRLR is fused with a GST tag. The Prolactin Receptor is purified by proprietary chromatographic techniques.
Species	Human
Source	E.coli
Protein Length	432-623 a.a.
Description	Prolactin is a pituitary hormone involved in the stimulation of milk production, salt and water regulation, growth, development and reproduction. The initial step in its action is the binding to a specific membrane receptor (prolactin receptor) which belongs to the superfamily of class 1 cytokine receptors. The function of the prolactin receptor is mediated, at least in part, by two families of signaling molecules: Janus kinases and signal transducers and activators of transcription. Prolactin (PRL) is a hormone involved in a variety of important functions including ion transport and osmoregulation, stimulation of milk, protein synthesis as well as the regulation of numerous reproductive functions. PRL exerts its influence on different cell types through a signal transduction pathway which begins with the binding of the hormone to a transmembrane PRL receptor. Immunoreactive PRL receptor, a member of the cytokine receptor family, varies in size (short and long forms) with tissue source and species, from ~40 kDa to 100 kDa. The PRL receptor consists of at least three

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

separate domains: an extracellular region with 5 cysteines which contains the prolactin binding site, a single transmembrane domain and a cytoplasmic region, the length of which appears to influence ligand binding and regulate cellular function.

Physical Appearance Sterile filtered liquid.

Formulation 100µpurified Human PRLR at 0.1 mg/ml in 50 mM Tris-Acetate, pH-7.5, 1 mM EDTA and 20% Glycerol.

Application And Suggested Dilutions • ELISA • Inhibition Assays • Western Blotting.

Biological Activity Not known.

Storage Store vial at -20°C to -80°C. When stored at the recommended temperature, this protein is stable for 12 months. Please avoid freeze-thaw cycles.

GENE INFORMATION

Gene Name PRLR prolactin receptor [Homo sapiens]

Synonyms PRLR; prolactin receptor; hPRLr; hPRL receptor; prolactin receptor delta 7/11; secreted prolactin binding protein; delta 4-SF1b truncated prolactin receptor; delta 4-delta 7/11 truncated prolactin receptor; OTTHUMP00000115998; PRL-R; prolactin receptor delta 7/11

Gene ID 5618

mRNA Refseq NM_000949

Protein Refseq NP_000940

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	176761
UniProt ID	P16471
Chromosome Location	5p13.2
Pathway	Cytokine-cytokine receptor interaction; Neuroactive ligand-receptor interaction; Jak-STAT signaling pathway
Function	ornithine decarboxylase activator activity; peptide hormone binding; prolactin receptor activity; protein binding; protein homodimerization activity; receptor activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA