

Recombinant Rainbow Trout Prolactin Receptor

Cat. No. prlr-61R Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Rainbow Trout Prolactin Receptor Extra Cellular Domain produced in <i>E. Coli</i> is a non-glycosylated, Polypeptide chain containing 210 amino acids and having a molecular mass of 24034 Dalton. The Prolactin Receptor is purified by proprietary chromatographic techniques.
Species	Rainbow Trout
Source	E.coli
Description	Prolactin is a pituitary hormone involved in the stimulation of milk production, salt and water regulation, growth, development and reproduction. The initial step in its action is the binding to a specific membrane receptor (prolactin receptor) which belongs to the superfamily of class 1 cytokine receptors. The function of the prolactin receptor is mediated, at least in part, by two families of signaling molecules: Janus kinases and signal transducers and activators of transcription. Prolactin (PRL) is a hormone involved in a variety of important functions including ion transport and osmoregulation, stimulation of milk, protein synthesis as well as the regulation of numerous reproductive functions. PRL exerts its influence on different cell types through a signal transduction pathway which begins with the binding of the hormone to a transmembrane PRL receptor. Immunoreactive PRL receptor, a member of the cytokine receptor family, varies in size (short and long forms) with tissue source and species, from ~40 kDa to 100 kDa. The PRL receptor consists of at least three separate domains: an extracellular region with 5 cysteines which contains the prolactin binding site, a single transmembrane domain and a cytoplasmic region, the

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	length of which appears to influence ligand binding and regulate cellular function.
Physical Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Purity	Greater than 97.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.
Formulation	The protein was lyophilized from a concentrated (1mg/ml) solution with 0.0045mM NaHCO ₃ .
Solubility	It is recommended to reconstitute the lyophilized PRL-R in sterile 18MΩ-cm H ₂ O not less than 100g/ml, which can then be further diluted to other aqueous solutions.
Amino Acid Sequence	The sequence of the first five N-terminal amino acids was determined and was found to be Ala-Arg-His-Thr-Pro.
Biological Activity	Activity is determined by the dose-dependant inhibition of Prolactin-stimulated proliferation of Nb2 cells and by high affinity binding of oPLR and other lactogenic hormones.
Protein Content	Protein quantitation was carried out by two independent methods: 1.UV spectroscopy at 280 nm using the absorbency value of 2.48 as the extinction coefficient for a 0.1% (1mg/ml) solution. This value is calculated by the PC GENE computer analysis program of protein sequences (IntelliGenetics). 2. Analysis by RP-HPLC, using a standard solution of PRLr-ECD as a Reference Standard.
Storage	Lyophilized PRL-R although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Prolactin Receptor should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent

freeze-thaw cycles.

GENE INFORMATION

Gene Name [prlr prolactin receptor \[Oncorhynchus mykiss \]](#)

Synonyms prolactin receptor; prlr

Gene ID [100136609](#)

mRNA Refseq [NM_001124599](#)

Protein Refseq [NP_001118071](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA