

Recombinant Human IL9 protein

Cat. No. IL9-67H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human IL9 protein was expressed in Escherichia coli.
Species	Human
Source	E.coli
ProteinLength	126
Description	Interleukin-9 (IL-9) is encoded by the IL9 gene and produced by T-cells and specifically by CD4+ helper cells. IL-9 was originally identified as a cytokine found in the conditioned medium of a human T cell leukemia virus type I (HTLVI) transformed T cell line. It functions through the IL-9 receptor, which activates different signal transducer and activator (STAT) proteins and thus connects this cytokine to various biological processes. IL-9 can support the growth of IL-2 independent and IL-4 independent helper T-cells. Human IL-9 has approximately 56 % amino acid sequence identity with murine IL-9. The gene encoding this cytokine has been identified as a candidate gene for asthma. Genetic studies on a mouse model of asthma demonstrated that this cytokine is a determining factor in the pathogenesis of bronchial hyper-responsiveness.
Form	Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by a cell proliferation assay using human MO7e cells is less than 0.2 ng/ml, corresponding to a specific activity of > 5.0 ×

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	10 ⁶ IU/mg.
Molecular Mass	Approximately 14.1 kDa, a single non-glycosylated polypeptide chain containing 126 amino acids.
AA Sequence	QGCPTLAGILDINFLINKMQEDPASKCHCSANVTSCCLCLGIPSDNCTRPCFSERLSQ MTNTTMQTRYPLIFSRVKKSVEVLKNNKCPYFSCEQPCNQTTAGNALTFLKSLL EIF QKEKMRGMRGKI
Endotoxin	Less than 1 EU/g of rHuIL-9 as determined by LAL method.
Purity	>95% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	IL9
Official Symbol	IL9
Synonyms	IL9; interleukin 9; interleukin-9; homolog of mouse T cell and mast cell growth factor

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

40; HP40; IL 9; P40; p40 cytokine; p40 T cell and mast cell growth factor; T cell growth factor p40; cytokine P40; T-cell growth factor p40; p40 T-cell and mast cell growth factor; IL-9;

Gene ID [3578](#)


mRNA Refseq [NM_000590](#)

Protein Refseq [NP_000581](#)

MIM [146931](#)

UniProt ID [P15248](#)

SDS-PAGE of IL9-67H



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA