

Recombinant Mouse Interleukin 15

Cat. No. IL15-69M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Mouse Interleukin-15 produced in <i>E. Coli</i> is a single, non-glycosylated polypeptide chain containing 115 amino acids and having a molecular mass of 13321 Dalton. The IL-15 is purified by proprietary chromatographic techniques.
Species	Mouse
Source	E.coli
Description	The protein encoded by this gene is a cytokine that regulates T and natural killer cell activation and proliferation. This cytokine and interleukine 2 share many biological activities. They are found to bind common hematopoietin receptor subunits, and may compete for the same receptor, and thus negatively regulate each other's activity. The number of CD8+ memory cells is shown to be controlled by a balance between this cytokine and IL2. This cytokine induces the activation of JAK kinases, as well as the phosphorylation and activation of transcription activators STAT3, STAT5, and STAT6. Studies of the mouse counterpart suggested that this cytokine may increase the expression of apoptosis inhibitor BCL2L1/BCL-x(L), possibly through the transcription activation activity of STAT6, and thus prevent apoptosis. Two alternatively spliced transcript variants of this gene encoding the same protein have been reported.
Amino Acid Sequence	The sequence of the first five N-terminal amino acids was determined and was found to be Met-Asn-Trp-Ile-Asp.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Physical Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Purity	Greater than 98% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.
Formulation	The protein was lyophilized from a concentrated (1mg/ml) solution with 10mM NaP pH 8.0.
Solubility	It is recommended to reconstitute the lyophilized Interleukin-15 in sterile 18MΩ-cm H ₂ O not less than 100g/ml, which can then be further diluted to other aqueous solutions.
Biological Activity	The ED ₅₀ as determined by the dose-dependant stimulation of the proliferation of mouse CTLL-2 was found to be 15 ng/ml, corresponding to a Specific Activity of 2 x 10 ⁵ IU/mg.
Storage	Lyophilized Interleukin-15 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL15 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles

GENE INFORMATION

Gene Name	IL15 interleukin 15 [<i>Mus musculus</i>]
Synonyms	IL15; interleukin 15; AI503618
Gene ID	16168

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA Refseq	NM_008357
Protein Refseq	NP_032383
UniProt ID	P48346
Chromosome Location	8 C2; 8 38.0 cM
Pathway	Cytokine-cytokine receptor interaction; Jak-STAT signaling pathway
Function	cytokine activity; cytokine receptor binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA