

Recombinant Human Regulator Of Calcineurin 2, His-tagged

Cat. No. RCAN2-4750H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human RCAN2 protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.
Species	Human
Source	Human
Description	Calcipressin-2, also known as RCAN2, belonging to the RCAN (Regulator of calcineurin) family modulate the activity of Calcineurin through a valine-rich region within their carboxyl tail. RCANs compete with NFAT(nuclear factor of activated T-cells) for binding to the same docking site in calcineurin, which results in either inhibition or activation of calcineurin activity. RCAN2 is a Ca(2+) dependent protein phosphatase that is involed in several neuornal functions.
Form	Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 1mM DTT, 20% glycerol, 0.2M NaCl
Molecular Weight	24.1 kDa (217aa), confirmed by MALDI-TOF
Purity	> 85% by SDS - PAGE
Concentration	0.5 mg/ml (determined by Bradford assay)
Sequences of amino acids	MGSSHHHHHH SSGLVPRGSH MPAPSMDCDV STLVACVVDV EVFTNQEVKE KFEGLFRTYD DCVTFQLFKS FRRVRINFSN PKSAARARIE LHETQFRGKK

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

LKLYFAQVQT PETDGDKLHL APPQPAKQFL ISPPSSPPVG WQPINDATPV
 LNYDLLYAVA KLGPGEKYEL HAGTESTPSV VVHVCDSDIE EEEDPKTSPK
 PKIIQTRRPG LPPSVSN

Storage

Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

Pathways

Calcium signaling in the CD4+ TCR pathway; Role of Calcineurin-dependent NFAT signaling in lymphocytes

GENE INFORMATION

Gene Name

[RCAN2 regulator of calcineurin 2 \[Homo sapiens \]](#)

Official Symbol

[RCAN2](#)

Synonyms

RCAN2; CSP2; RCN2; MCIP2; ZAKI4; hRCN2; ZAKI-4; DSCR1L1; regulator of calcineurin 2; calciopressin-2; OTTHUMP00000016544; Down syndrome candidate region 1-like 1; thyroid hormone-responsive protein ZAKI-4; Down syndrome critical region gene 1-like 1; thyroid hormone-responsive (skin fibroblasts); myocyte-enriched calcineurin-interacting protein 2; Calciopressin 2 (Thyroid hormone-responsive protein ZAKI-4) (Down syndrome candidate region 1-like 1) (Myocyte-enriched calcineurin interacting protein 2) (MCIP2)

Gene ID

[10231](#)

mRNA Refseq

[NM_005822](#)

Protein Refseq

[NP_005813](#)

MIM

[604876](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	Q14206
Chromosome Location	6p12.3
Function	nucleotide binding; protein phosphatase 2B binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA