

Active Recombinant Mouse Tumor Necrosis Factor (Ligand) Superfamily, Member 11

Cat. No. Tnfsf11-474M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant M-sTnfsf11 is approximately 19.9 kDa and contains 176 amino acids.
Species	Mouse
Source	E.coli
Description	Tnfsf11 occurs in 3 forms: cell-bound Tnfsf11, which is expressed by osteoblast lineage cells, soluble Tnfsf11, which is expressed by activated T lymphocytes, and a truncated ectodomain form derived from the cell-bound Tnfsf11, which is enzymatically processed by TACE. All three forms stimulate their specific receptor, RANK, which is located on osteoclastic and dendritic cells. Tnfsf11 is critically involved in osteoclastic function, and in the regulation of specific immunity.
Form	Lyophilized 10 mM NaP, pH 7.5 + 50 mM NaCl.
Purity	Purity Greater than 95% as determined by: Analysis by RP-HPLC. Reducing and non-reducing SDS-PAGE.
Bio-activity	Measured by its ability to induce osteoclast formation on mouse RAW264.7 cells using a concentration of 5.0-10.0 ng/ml.
Physical Appearance	Sterile Filtered white lyophilized (freeze-dried) powder.
Solubility	Centrifuge vial before opening. When reconstituting the product, gently pipet and

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at a concentration of 0.1 mg/μs/ml, which can be further diluted into other aqueous solutions.

Purity Greater than 95% as determined by

Analysis by RP-HPLC. Reducing and non-reducing SDS-PAGE.

Protein Content determined by

UV spectroscopy at 280 nm. Analysis by RP-HPLC calibrated against a known standard. Quantitation on SDS-PAGE against a known standard.

Endotoxin Level

Endotoxin level as measured by LAL is <0.01ng/ug or <0.1EU/ug.

Amino acid sequence

PAMMEGSWLD VAQRGKPEAQ PFAHLTINAA SIPSGSHKVT LSSWYHDRGW
AKISNMTLSN GKLRVNQDGF YYLYANICFR HHETSGSVPT DYQLQMVYVV
KTSIKIPSSH NLMKGGSTKN WSGNSEFHFY SINVGGF FKL RAGEEISIQV
SNPSLLDPDQ DATYFGAFKV QDID

Storage

Lyophilized product is very stable at -20 degrees. Reconstituted material should be aliquoted and frozen at -20. It is recommended to add a carrier protein (0.1% HSA or BSA) for long term storage.

GENE INFORMATION

Gene Name

Tnfsf11 tumor necrosis factor (ligand) superfamily, member 11 [*Mus musculus*]

Official Symbol

Tnfsf11

Synonyms

Tnfsf11; tumor necrosis factor (ligand) superfamily, member 11; ODF; OPG; OPGL; RANKL; Ly109l; Trance; OPG ligand; osteoprotegerin ligand; osteoclast

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

differentiation factor; receptor activator of NF-kappaB ligand; tumor necrosis factor-related activation-induced cytokine; Ly109I

Gene ID [21943](#)

mRNA Refseq [NM_011613](#)

Protein Refseq [NP_035743](#)

UniProt ID [O35235](#)

Chromosome Location 14 D3; 14 45.0 cM

Pathway Cytokine-cytokine receptor interaction

Function cytokine activity; protein binding; receptor activity; tumor necrosis factor receptor binding

PDB rendering based on 1s55

