

Recombinant Vascular Endothelial Growth Factor-E (Orf Virus)

Cat. No. VEGFE-52 **Lot. No.** (See product label)

SPECIFICATION

Product Overview

A DNA sequence encoding the mature variant of ov VEGF-E isolate D1701 (Dehio et al., 1999; GenBank accession No. AF106020) was expressed in E. coli as a 132 amino acid residue fusion protein with an N-terminal His-tag sequence and a thrombin cleavage site. Recombinant VEGF-E homodimer was dimerized in vitro and has a predicted mass of approximately 35 kDa. Based on sequence similarity to VEGF-A, a gene encoding a VEGF homologue has recently been discovered in the genome of Orf virus (OV) (Lyttle et al., 1994). Different isolates of Orf virus show significant amino acid sequence similarity to VEGF-A and described as a viral virulence factor that appears to be derived from captured host genes. All eight cysteine residues of the central cysteine knot motif characteristic of members of the VEGF family are conserved among other residues in the VEGF-E proteins (Dehio et al., 1999; Wise et al., 1999). Alignment of all mammalian VEGF sequences indicated that VEGF-E is distinct from the previously described VEGFs but most closely related to VEGF-A. Like VEGF-A, VEGF-E was found to bind with high affinity to VEGF receptor-2 (KDR) resulting in receptor autophosphorylation, whilst in contrast to VEGF-A, VEGF-E can not bind to VEGF receptor-1 (Flt-1). Furthermore VEGF-E can also not bind to VEGF receptor-3 (FLT-4). Therefore VEGF-E is a potent angiogenic factor selectively binding to VEGF receptor –2/KDR.

Source

E.coli

Physical Appearance

Sterile Filtered White lyophilized (freeze-dried) powder.

Purity

Greater than 90.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	SDS-PAGE.
Formulation	The protein was lyophilized from a concentrated (1mg/ml) solution containing PBS pH-7.4.
Solubility	The lyophilized VEGF-E is soluble in water and most aqueous buffers. The lyophilized Vascular Endothelial Growth Factor-E (Orf Virus) should be reconstituted in PBS or medium containing at least 0.1% human or bovine serum albumin to a concentration not lower than 50g/ml.
Biological Activity	The ED ₅₀ for stimulation of 3H-thymidine incorporation and cell proliferation by human umbilical vein endothelial cells for VEGF-E Orf Virus has been determined to be in the range of 5 - 20 ng/ml, corresponding to a specific activity of 2 x 10 ⁵ Units/mg.
Storage	Lyophilized Vascular Endothelial Growth Factor-E Orf Virus although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution VEGF E -OV should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.

GENE INFORMATION

Synonyms	VEGF-E; Vascular Endothelial Growth Factor-E.
-----------------	---

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA