

Recombinant Rat Tnf protein

Cat. No. Tnf-71R **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Rat Tnf protein was expressed in Escherichia coli.
Species	Rat
Source	E.coli
ProteinLength	156
Description	Tumor necrosis factor alpha (TNF-α), also called cachectin, is the best-known member of the TNF-family, which can cause cell death. This protein is produced by neutrophils, activated lymphocytes, macrophages, NK cells, LAK cells, astrocytes, endothelial cells, smooth muscle cells and some transformed cells. TNF-α occurs as a secreted, soluble form and as a membrane-anchored form, both of which are biologically active. The naturally-occurring form of TNF-α is glycosylated, but non-glycosylated recombinant TNF-α has comparable biological activity. Two types of receptors for TNF-α have been described and virtually all cell types studied show the presence of one or both of these receptor types.
Form	Lyophilized from a 0.2 μ m filtered concentrated solution in 20 mM PB, pH 7.2, 150 mM NaCl.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by a cytotoxicity assay using murine L929 cells is less than 0.05 ng/ml, corresponding to a specific activity of $> 2.0 \times 10^7$ IU/mg in the presence of actinomycin D.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	Approximately 17.2 kDa, a single non-glycosylated polypeptide chain containing 156 amino acids.
AA Sequence	LRSSSQNSSDKPVAHVVANHQAEEQLEWLSQRANALLANGMDLKDNLVVPADGL YLIYSQVLFGKGQCPDYVLLTHTVSRFAISYQEKVSLLSAIKSPCPKDTPEGAELKPW YEPMYLGGVFQLEKGDLLSAEVNLPKYLDITESGQVYFGVIAL
Endotoxin	Less than 1 EU/μg of rRtTNF-α as determined by LAL method.
Purity	>98% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	Tnf
Official Symbol	Tnf
Synonyms	Tumor Necrosis Factor, TNFSF2, Cachectin, Differentiation-inducing Factor , DIF, Necrosin, Cytotoxin

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


Gene ID 24835

mRNA Refseq NM_012675

Protein Refseq NP_036807

UniProt ID P16599

SDS-PAGE of Tnf-71R



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA