

Recombinant Human TXNDC12, His-tagged

Cat. No. TXNDC12-3498H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human TXNDC12(His 27 - Leu 172), fused with 6×His tag at the C-terminus, was expressed in human 293 cells.
Species	Human
Source	HEK293
ProteinLength	27-172 a.a.
Description	This gene encodes a member of the thioredoxin superfamily. Members of this family are characterized by a conserved active motif called the thioredoxin fold that catalyzes disulfide bond formation and isomerization. This protein localizes to the endoplasmic reticulum and has a single atypical active motif. The encoded protein is mainly involved in catalyzing native disulfide bond formation and displays activity similar to protein-disulfide isomerases. This protein may play a role in defense against endoplasmic reticulum stress. Alternate splicing results in both coding and non-coding variants.
Form	Lyophilized from 0.22 µm filtered solution in PBS, pH7.4. Normally Mannitol or Trehalose are added as protectants before lyophilization.
Molecular Mass	rhTXNDC12, fused with 6×His tag at the C-terminus, has a calculated MW of 17.3 kDa. The predicted N-terminus is His 27. DTT-reduced Protein migrates as 18-21 kDa.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	Less than 1.0 EU per µg of the rhTXNDC12 by the LAL method.
Purity	>95% as determined by SDS-PAGE.
Storage	Avoid repeated freeze-thaw cycles. No activity loss was observed after storage at: In lyophilized state for 1 year (4°C); After reconstitution under sterile conditions for 3 months (-70°C).

GENE INFORMATION

Gene Name	TXNDC12 thioredoxin domain containing 12 (endoplasmic reticulum) [Homo sapiens (human)]
Official Symbol	TXNDC12
Synonyms	TXNDC12; thioredoxin domain containing 12 (endoplasmic reticulum); thioredoxin domain-containing protein 12; AGR1; anterior gradient homolog 1 (Xenopus laevis); endoplasmic reticulum thioredoxin superfamily member; 18 kDa; ERP18; ERP19; hAG 1; PDIA16; protein disulfide isomerase family A; member 16; TLP19; ER protein 18; ER protein 19; anterior gradient homolog 1; thioredoxin-like protein p19; endoplasmic reticulum protein ERp19; endoplasmic reticulum resident protein 18; endoplasmic reticulum resident protein 19; protein disulfide isomerase family A, member 16; endoplasmic reticulum thioredoxin superfamily member, 18 kDa; AG1; ERP16; hAG-1; hTLP19
Gene ID	51060
mRNA Refseq	NM_015913
Protein Refseq	NP_056997

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	609448
UniProt ID	O95881
Chromosome Location	1p32.3
Pathway	Glutathione metabolism
Function	protein binding; protein-disulfide reductase (glutathione) activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA