

Recombinant Human, HSPA5 His-tagged

Cat. No. HSPA5-3677H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Total 646 AA. MW: 71.72 kDa (calculated). N-Terminal His-tag, 10 extra AA (highlighted).
Species	Human
Source	E.coli
Description	Binding immunoglobulin protein (BiP) also known as 78 kDa glucose-regulated protein (GRP-78) or heat shock 70 kDa protein 5 (HSPA5) is a HSP70 molecular chaperone located in the lumen of the endoplasmic reticulum (ER), that binds newly-synthesized proteins as they are translocated into the ER and maintains them in a state competent for subsequent folding and oligomerization. BiP is also an essential component of the translocation machinery, as well as playing a role in retrograde transport across the ER membrane of aberrant proteins destined for degradation by the proteasome. BiP is an abundant protein under all growth conditions, but its synthesis is markedly induced under conditions that lead to the accumulation of unfolded polypeptides in the ER. In humans, BiP is encoded by the HSPA5 gene.
Form	Filtered (0,4 µm) and lyophilized in 0.5 mg/mL in 20mM TRIS, 50mM NaCl, pH 7.5
Reconstitution	Add deionized water to prepare a working stock solution of approximately 0.5 mg/mL and let the lyophilized pellet dissolve completely. Product is not sterile! Please filter the product by an appropriate sterile filter before using it in the cell culture.
Applications	Western blotting

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Quality Control Test	BCA to determine quantity of the protein; SDS PAGE to determine purity of the protein.
Sequences of amino acids	MKHHHHHHAS EEEDKKEDVG TVVGIDLGTT YSCVGVFKNG RVEIANDQG NRITPSYVAF TPEGERLIGD AAKNQLTSNP ENTVFDAKRL IGRTWNDPSV QQDIKFLPFK VVEKKTkPYI QVDIGGGQTK TFAPEEISAM VLTKMKETAE AYLGKKVTHA VVTVPAYFND AQRQATKDAG TIAGLNMRI INEPTAAAIA YGLDKREGEK NILVFDLGGG TFDVSLLTID NGVFEVVATN GDTHLGGEDF DQRMMEHFik LYKKKTGKDV RKDNRAVQKL RREVEKAKRA LSSQHqARIE IESFYEGEDF SETLTRAKFE ELNMDLFRST MKPVQKVLED SDLKKSIDIE IVLVGGSTRI PKIQQLVKEF FNGKEPSRGI NPDEAVAYGA AVQAGVLSGD QDTGDLVLLD VCPLTLGIET VGGVMTKLIP RNTVVPTKKS QIFSTASDNQ PTVTIKVYEG ERPLTKDNHL LGTFDLTGIP PAPRGVPQIE VTFEIDVNGI LRVTAEDKGT GNKNKITITN DQNRLTPEEI ERMVNDAEFK AEEDKKLKER IDTRNELESY AYSLKNQIGD KEKLGGKLSS EDKETMEKAV EEKIEWLESH QDADIEDFKA KKKELEEIVQ PIISKLYGSA GPPPTGEEDT AEKDEL
Storage	Store lyophilized protein at -20°C. Lyophilized protein remains stable until the expiry date when stored at -20°C. Aliquo treconstituted protein to avoid repeated freezing/thawing cycles and store at -80°C for long term storage. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after one week at 4°C.
GENE INFORMATION	
Gene Name	HSPA5 heat shock 70kDa protein 5 (glucose-regulated protein, 78kDa) [Homo sapiens]
Official Symbol	HSPA5

Synonyms	HSPA5; heat shock 70kDa protein 5 (glucose-regulated protein, 78kDa); HSPA5 heat shock 70kDa protein 5 (glucose-regulated protein, 78kDa); BIP; MIF2; GRP78; GRP 78; FLJ26106; BiP; OTTHUMP00000022124; Heat shock 70 kDa protein 5; Endoplasmic reticulum luminal Ca(2+)-binding protein grp78; Heat-shock 70kD protein-5 (glucose-regulated protein, 78kD); Immunoglobulin heavy chain-binding protein; heat shock 70kD protein 5 (glucose-regulated protein, 78kD); heat shock 70kDa protein 5
Gene ID	3309
mRNA Refseq	NM_005347
Protein Refseq	NP_005338
MIM	138120
UniProt ID	P11021
Chromosome Location	9q33-q34.1
Pathway	Antigen processing and presentation; Neurodegenerative Disorders; Prion disease; Hemostasis.
Function	ATP binding; calcium ion binding; caspase inhibitor activity; nucleotide binding; protein binding, bridging; ribosome binding; unfolded protein binding.