

Active Recombinant Human CD86 Protein, Fc Chimera

Cat. No. CD86-575H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human CD86 (Accession # NP_008820), fused with , was produced in Chinese Hamster Ovary cell line, CHO-derived.
Species	Human
Source	CHO
Predicted N Terminal	Leu20
Form	Lyophilized from a 0.2 µ filtered solution in PBS.
Bio-activity	Measured by its ability to induce IL-2 secretion by Jurkat human acute T cell leukemia cells. Freeman, G.J. et al. (1993) Science 262:909. The ED50 for this effect is typically 0.04-0.2 µg/mL.
Molecular Mass	Recombinant Human CD86, Fc Chimera has a calculated MW of 52 kDa (monomer). In SDS-PAGE migrates as 75-100 kDa, reducing conditions.
Purity	>95%, by SDS-PAGE under reducing conditions and visualized by silver stain.
Storage	Avoid repeated freeze-thaw cycles. No activity loss was observed after storage at: In lyophilized state for 1 year (4°C); After reconstitution under sterile conditions for 3 months (-70°C).
Reconstitution	Reconstitute at 250 µg/mL in PBS containing at least 0.1% human or bovine serum

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

albumin.

GENE INFORMATION

Gene Name [CD86 CD86 molecule \[Homo sapiens \]](#)

Official Symbol CD86

Synonyms

CD86; CD86 molecule; CD28LG2, CD86 antigen (CD28 antigen ligand 2, B7 2 antigen); T-lymphocyte activation antigen CD86; B lymphocyte antigen B7 2; B7 2; B7.2; BU63; FUN-1; CTLA-4 counter-receptor B7.2; B-lymphocyte activation antigen B7-2; CD86 antigen (CD28 antigen ligand 2, B7-2 antigen); B70; B7-2; LAB72; CD28LG2; MGC34413;

Gene ID [942](#)

mRNA Refseq [NM_001206924](#)

Protein Refseq [NP_001193853](#)

MIM [601020](#)

UniProt ID P42081

Chromosome Location 3q21

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA