

Active Recombinant Human TNFSF15 protein

Cat. No. TNFSF15-1071H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human TNFSF15 protein was expressed in Escherichia coli.
Species	Human
Source	E.coli
ProteinLength	72-251 a.a.
Description	The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This protein is abundantly expressed in endothelial cells, but is not expressed in either B or T cells. The expression of this protein is inducible by TNF and IL-1 alpha. This cytokine is a ligand for receptor TNFRSF25 and decoy receptor TNFRSF21/DR6. It can activate NF-kappaB and MAP kinases, and acts as an autocrine factor to induce apoptosis in endothelial cells. This cytokine is also found to inhibit endothelial cell proliferation, and thus may function as an angiogenesis inhibitor. Two transcript variants encoding different isoforms have been found for this gene.
Form	Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4, with 0.02% Tween-20.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by its ability to induce apoptosis using human TF-1 cells is less than 20 ng/ml, corresponding to a specific activity of $> 5.0 \times 10^4$ IU/mg.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	Approximately 20.5 kDa, a single non-glycosylated polypeptide chain containing 180 amino acids.
AA Sequence	LKGQEFAPSHQQVYAPLRADGDKPRAHLTVVRQTPTQHFKNQFPALHWEHELGLA FTKNRMNYTNKFLIPESGDYFIYSQVTFRGMTSECSEIRQAGRPNKPDSITVVITKV TDSYPEPTQLLMGTKSVCEVGSNWFQPIYLGAMFSLQEGDKLMVNVSDISLVDYTK EDKTFFGAFL
Endotoxin	Less than 0.1 EU/g of rHuTL-1A/TNFSF15 as determined by LAL method.
Purity	>97% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	TNFSF15
Official Symbol	TNFSF15
Synonyms	TNFSF15; tumor necrosis factor (ligand) superfamily, member 15; tumor necrosis factor ligand superfamily member 15; MGC129934; MGC129935; TL1; TL1A; TNF

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

ligand related molecule 1; TNF superfamily ligand TL1A; vascular endothelial cell growth inhibitor; vascular endothelial growth inhibitor 192A; VEGI; VEGI192A; TNF ligand-related molecule 1; vascular endothelial growth inhibitor-192A;

Gene ID 9966


mRNA Refseq NM_001204344

Protein Refseq NP_001191273

MIM 604052

UniProt ID O95150

**SDS-PAGE of
TNFSF15-1071H**



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA