

Lysozyme From Chicken Eggs (Food Grade)

Cat. No. LYZ-65E **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species Chicken

Source Chicken Egg

Description

Lysozyme is an enzyme found in egg white, tears, and other secretions. Lysozyme is responsible for attacking & breaking down the polysaccharide walls of many kinds of bacteria and thus it provides some protection against infection. Bacteria build a tough skin of carbohydrate chains, interlocked by short peptide strands, that braces their delicate membrane against the cell's high osmotic pressure. Lysozyme breaks these carbohydrate chains, destroying the structural integrity of the cell wall. The bacteria burst under their own internal pressure. Lysozyme (muramidase) hydrolyzes preferentially the β -1,4 glucosidic linkages between N-acetylmuramic acid and N-acetylglucosamine which occur in the mucopeptide cell wall. Lysozyme is a mediator in the anti-tumor function of macrophages which, it has been shown, secrete the enzyme. Cartilage lysozyme has a role in cartilage calcification. Presence of the Lysozyme in cerebrospinal fluid is indicative of tumor of the central nervous system, normally lysozyme activity is practically absent from urine, bile and spinal fluid.

Appearance Sterile Filtered white lyophilized powder.

CAS Number 12650-88-3

Specific Activity $\geq 20\text{KU/mg protein}$

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Unit definition	One unit will produce a change in A450 of 0.001 per minute at pH 6.24 at 25 oC, using a suspension of <i>Micrococcus lysodeikticus</i> as substrate, in a 2.6 ml reaction mixture (1 cm light path).
Molecular Weight	14.3 kDa (amino acid sequence)
Isoelectric point	11.35
Optimum temperature	2-8 oC
Optimum pH	5.3-6.4
Inhibitors	Lysozyme is inhibited by indole derivatives, which bind to and distort the active site, and imidazole, which induces the formation of a charge-transfer complex. It is also inhibited by surface-active agents such as sodium dodecyl sulfate, sodium dodecanate, and dodecyl alcohol. Other compounds of these types with carbon chains of 12 or more carbons in length will also inhibit lysozyme.
Preparation Instructions	For E. coli cell lysis, use a freshly prepared lysozyme solution (10 mg/ml) in 10 mM Tris-HCl, pH 8.0. Aqueous solutions should retain activity for at least one month when stored between 2–8 oC.
Storage Temperature	2-8 oC
GENE INFORMATION	
Gene Name	LYZ lysozyme (renal amyloidosis) [<i>Gallus gallus</i>]
Official Symbol	LYZ

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Synonyms	LYZ; lysozyme; LYZC
Gene ID	396218
mRNA Refseq	NM_205281
Protein Refseq	NP_990612
UniProt ID	P00698
Function	hydrolase activity, acting on glycosyl bonds; lysozyme activity; protein binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA