

LL/2 (mouse Lewis lung carcinoma) whole cell lysate

Cat. No. LL-2-991M **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species	Mouse
Tissue Type	Lung
Preparation method	Cell lysate was prepared by homogenization in ice-cold modified RIPA Lysis Buffer with cocktail of protease inhibitors. Cell debris was removed by centrifugation. Protein concentration was determined with Bradford assay (Bio-Rad protein assay, Microplate Standard assay). The lysate was adjusted to 2 mg/ml.
Lysis buffer	Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF.
Size	200 ug
Recommended Usage	Use it directly for immuno-precipitation, or heat lysate with SDS gel loading buffer to 95°C for 5 minutes followed by rapid cooling for western blot application. If dissociating conditions are required, add reducing agent prior to heating.
Storage Buffer	In modified RIPA Lysis Buffer.
Storage Instruction	Store at -80°C. Aliquot to avoid repeated freezing and thawing.
Applications	Western Blot;Immunoprecipitation;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA