

Nickel Beads

Cat. No. Nickel-31A **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Recombinant proteins tagged with 6-10 poly Histidines can be purified in one step by ion metal affinity chromatography (IMAC). Several metal ions exhibiting affinity to poly-Histidines are commonly used, Nickel and Cobalt being the most popular. Tagging the proteins may be done at the N-ter or the C-ter by cloning a gene in dedicated expression vectors encoding to the poly Histidines stretch, enabling the purification of the recombinant protein resulted.

Cat.no Nickel-31A

Matrix Highly cross-linked spherical agarose, 4%

Average particle size 40-165 µm

Dynamic binding capacity ~6-9 mg pure (His) 6 -tagged protein (Mr ~ 49 000) per ml

Recommended flow rate 1-2 ml/min/ cm2

pH working range 2-11

Elution buffer 0.02 M sodium phosphate, 0.5 M NaCl, 0.5 M imidazole, pH 7.4

Storage Storage conditions: equilibrate the column with the Binding buffer added with NaN3 0.1% (w/v) or 20% ethanol as preservatives.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA