

Active Recombinant Human TEK, Fc-tagged, Biotinylated

Cat. No. TEK-702H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	The recombinant human Tie2-Fc fusion is expressed as a 952 amino acid protein consisting of Ala23- Met746 region of Tie2 and a C-terminal Fc from human IgG1, which exists as a dimer under non-reducing conditions.
Species	Human
Source	Human Cells
Form	Supplied at 0.5 mg/ml in sterile PBS pH7.4 (carrier & preservative free). The purified recombinant protein was labeled with Biotin (3-5 Biotin per molecule).
Bio-activity	Recombinant Tie2 protein binds human Angiopoietin-2 and inhibits Angiopoietin-2 mediated signaling activity.
Molecular Mass	Calculated molecular mass (kDa): 106.3; Estimated by SDS-PAGE under reducing condition (kDa): 110-120
AA Sequence	AMDLILINSLPLVSDAETSLTCIASGWRPHEPITIGRDFEALMNQHQPDPLEVTQDVTR EWAKKVVWKREKASKI NGAYFCEGRVRGEAIRIRTMKMRQQASFLPATLTMTVDK GDNVNISFKKVLIKEEDAVIYKNGSFIHSVPRHEV PDILEVHLPHAQPQDAGVYSARY IGGNLFTSAFTRLIVRRCEAQKWGPECNHLCTACMNNGVCHEDTGECICPP GFMG RTCEKACELHTFGRTCKERCSGQEGCKSYVFCLPDYPYGCSCATGWKGLQCNEAC HPGFYGPDCCKLRCSN NGEMCDRFQGCCLSPGWQGLQCEREGIPRMTPKIVDLP DHIEVN SGKFN PICKASGWPLPTNEEMTLVKPDGTV LHPKDFNHTDHFSVAIFTIHR LPPDSGVVWCSVNTVAGMVEKPFNISVKVLPKPLNAPNVIDTGHNFAVINI SSEPYF

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

GDGPIKSKLLYKPVNHYEAWQHIQVTNEIVTLNYLEPRTEYELCVQLVRRGEGGEG
 HPGPVRRFTTA SIGLPPPRGLNLLPKSQTTLNLTWQPIFPSSDDFYVEVERRSVQK
 SDQQNIKVPGNLTSVLLNNLHPREQYVV RARVNTKAQGEWSEDLTAWTSLDILPPQ
 PENIKISNITHSSAVISWTILDGYSISSITIRYKVQGKNEDQHVDV KIKNATITQYQLKG
 LEPETAYQVDIFAENNIGSSNPAFSHELVTLPESQAPADLGGGKMSTGHTCPCPA
 PEL LGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNA
 KTKPREEQYNSTYRVVSVLT VLNQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQP
 REPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEW ESNGQPENNYKTPPVLD
 SDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

Endotoxin

<0.1 eu="" per="" 1="" µg="" of="" purified="" recombinant="" protein=""
 determined="" by="" the="" la="">

Purity

>95% judged by SDS-PAGE under reducing condition

Storage

The product is shipped at 4°C. Upon receipt, centrifuge the product briefly before opening the vial. It is recommended to store small aliquots at the temperature below -20°C for long-term storage and the product is stable for 3 months. The undiluted protein can be stored at 4°C for no more than 2 weeks. Avoid repeated freeze-thaw cycles.

Conjugation

Biotin

GENE INFORMATION

Gene Name

TEK TEK tyrosine kinase, endothelial [Homo sapiens]

Official Symbol

TEK

Synonyms

TEK; TEK tyrosine kinase, endothelial; venous malformations, multiple cutaneous and mucosal , VMCM; angiopoietin-1 receptor; CD202b; TIE 2; TIE2; VMCM1; hTIE2;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	p140 TEK; soluble TIE2 variant 1; soluble TIE2 variant 2; endothelial tyrosine kinase; tyrosine-protein kinase receptor TEK; tunica interna endothelial cell kinase; tyrosine-protein kinase receptor TIE-2; tyrosine kinase with Ig and EGF homology domains-2; VMCM; TIE-2; CD202B;
Gene ID	7010
mRNA Refseq	NM_000459
Protein Refseq	NP_000450
MIM	600221
UniProt ID	Q02763
Chromosome Location	9p21
Pathway	Angiogenesis, organism-specific biosystem; Angiopoietin receptor Tie2-mediated signaling, organism-specific biosystem; Cell surface interactions at the vascular wall, organism-specific biosystem; Hemostasis, organism-specific biosystem; Rheumatoid arthritis, organism-specific biosystem; Rheumatoid arthritis, conserved biosystem; Tie2 Signaling, organism-specific biosystem;
Function	ATP binding; nucleotide binding; protein binding; protein kinase activity; protein tyrosine kinase activity; protein tyrosine kinase activity; receptor activity; transmembrane receptor protein tyrosine kinase activity;