

Recombinant Mouse IL7 protein

Cat. No. IL7-77M Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Mouse IL7 protein was expressed in Escherichia coli.
Species	Mouse
Source	E.coli
ProteinLength	129
Description	Interleukin-7 (IL-7) is encoded by the IL7 gene in mouse and secreted by stromal cells in the red marrow and thymus. The protein signals through the IL-7 receptor, which is a heterodimer consisting of IL-7 receptor alpha and IL-2 receptor gamma chain. IL-7 stimulates the differentiation of hematopoietic stem cells into lymphoid progenitor cells and it can stimulate proliferation of B cells, T cells and NK cells. Murine IL-7 has approximately 65 % and 88 % amino acid sequence identity with human and rat IL-7 and both proteins exhibit cross-species activity. IL-7 as an immunotherapy agent has been examined in many human clinical trials for various malignancies and during HIV infection.
Form	Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4, 2 % trehalose.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by a cell proliferation assay using murine 2E8 cells is less than 0.2 ng/ml, corresponding to a specific activity of $> 5.0 \times 10^6$ IU/mg.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	Approximately 14.9 kDa, a single non-glycosylated polypeptide chain containing 129 amino acids.
AA Sequence	ECHIKDKEGKAYESVLMISIDELDKMTGTDSNCPNNEPNFFRKHVCDDTKEAAFLNR AARKLKQFLKMNISEEFNVHLLTVSQGTQTLVNCTSKEEKNVKEQKKNDACFLKRLL REIKTCWNKILKGS
Endotoxin	Less than 1 EU/g of rMuIL-7 as determined by LAL method.
Purity	>96% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	IL7
Official Symbol	IL7
Synonyms	IL7; interleukin 7; interleukin-7; IL-7; hlb368; A630026I06Rik; MGC129342;
Gene ID	16196

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA Refseq NM_008371

Protein Refseq NP_032397

UniProt ID Q544C8

SDS-PAGE of II7-77M



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA