

Recombinant Human Fibroblast Growth Factor Receptor 2, Fc-tagged

Cat. No. FGFR2-02H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human Soluble FGFR-2a (IIIc) Fc Chimera fused with Xa cleavage site with the Fc part of human IgG1 produced in baculovirus is a heterodimeric, glycosylated, Polypeptide chain and having a molecular mass of 195 kDa. The FGFR2 is purified by proprietary chromatographic techniques.
Species	Human
Source	Insect Cells
Description	Fibroblast Growth Factors (FGFs) comprise a family of at least eighteen structurally related proteins that are involved in a multitude of physiological and pathological cellular processes, including cell growth, differentiation, angiogenesis, wound healing and tumorigenesis. The biological activities of the FGFs are mediated by a family of type I transmembrane tyrosine kinases which undergo dimerization and autophosphorylation after ligand binding. Four distinct genes encoding closely related FGF receptors, FGFR-1 to -4 are known. Multiple forms of FGFR-1 to -3 are generated by alternative splicing of the mRNAs. A frequent splicing event involving FGFR-1 and -2 results in receptors containing all three Ig domains, referred to as the alpha isoform, or only IgII and IgIII, referred to as the beta isoform. Only the alpha isoform has been identified for FGFR-3 and FGFR-4. Additional splicing events for FGFR-1 to -3, involving the C-terminal half of the IgIII domain encoded by two mutually exclusive alternative exons, generate FGF receptors with alternative IgIII domains (IIIb and IIIc). A IIIa isoform which is a secreted FGF binding protein

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

containing only the N-terminal half of the IgIII domain plus some intron sequences has also been reported for FGFR-1. Mutations in FGFR-1 to -3 have been found in patients with birth defects involving craniosynostosis.

Physical Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Purity	Greater than 90.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.
Formulation	CD332 was lyophilized from a concentrated (1 mg/ml) sterile solution containing no additives.
Solubility	It is recommended to reconstitute the lyophilized FGFR-2 in sterile PBS not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.
Biological Activity	Determined by its ability to inhibit human FGF acidic-dependent proliferation on R1 cells. The ED50 for this effect is typically at 15.0-30.0 ng/ml.
Storage	Lyophilized FGFR2A although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution FGFR2 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

GENE INFORMATION

Gene Name	FGFR2 fibroblast growth factor receptor 2 [Homo sapiens]
Synonyms	FGFR2; fibroblast growth factor receptor 2; BEK; JWS; CEK3; CFD1; ECT1; KGFR; TK14; TK25; BFR-1; CD332; K-SAM; FLJ98662; FGF receptor;

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

OTTHUMP00000020624; OTTHUMP00000020626; soluble FGFR4 variant 4; bacteria-expressed kinase; hydroxyaryl-protein kinase; keratinocyte growth factor receptor; BEK fibroblast growth factor receptor; protein tyrosine kinase, receptor like 14; EC 2.7.10.1; craniofacial dysostosis 1; Jackson-Weiss syndrome; Crouzon syndrome; Pfeiffer syndrome; FGFR-2;CD_antigen;CD332; Keratinocyte growth factor receptor 2; KSAM; OTTHUMP00000020621; OTTHUMP00000020629; FGF receptor

Gene ID	2263
mRNA Refseq	NM_000141
Protein Refseq	NP_000132
MIM	176943
UniProt ID	P21802
Chromosome Location	10q26
Pathway	Endocytosis; MAPK signaling pathway; Pathways in cancer; Prostate cancer; Regulation of actin cytoskeleton; Signaling by FGFR
Function	ATP binding; fibroblast growth factor receptor activity; heparin binding; nucleotide binding; protein binding; receptor activity; transferase activity

PDB rendering based
on 1djs.



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA