

Active Recombinant Mouse SERPINB2 protein, His-tagged

Cat. No. SERPINB2-14937M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Active Recombinant Mouse SERPINB2 protein(P12388)(1~415 aa), fused with N-terminal His tag, was expressed in HEK293.
Species	Mouse
Source	HEK293
ProteinLength	1-415 aa
Form	Lyophilized from sterile PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
Bio-activity	Serpin B2, also known as PAI-2, is an approximately 60 kDa serine protease inhibitor. It is primarily secreted by macrophages and monocytes and can form disulfide-linked multimers. Serpin B2 inhibits both the urokinase-type and tissue-type plasminogen activators (uPA and tPA). Serpin B2 also promotes the clearance of uPA by enhancing its binding and uptake by LRP. It limits fibril formation by Huntington protein (HTT) and beta-Amyloid peptides. It promotes Th2 biased immune responses and is important for intestinal CCL2 production, monocyte recruitment, and nematode clearance. A non-glycosylated form of Serpin B2 is retained intracellularly where it interferes with TNF-α induced apoptosis by protecting the Retinoblastoma protein (RB1) from calpain digestion. It also inhibits proteasome activity in activated endothelial cells. The activity of recombinant mouse PAI-2 was measured by its ability to inhibit uPA cleavage of a peptide substrate, N-carbobenzoyloxy-Gly-Gly-Arg-7-amido-4-methylcoumarin (Z-GGR-AMC)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

in the assay buffer 50 mM Tris, 0.01% Tween 20, pH 8.5. The 50 ul different concentrations of rmPAI-2 (MW: 47.9 KD) was incubated with 50ul 2ug/ml rhuPA (EPA140Mu 61) at room temperature for 15 minutes. Loading 50 μ of the incubated mixtures into empty wells of a plate, and start the reaction by adding 50 μ of 200 μM substrate (Z-GG R-AMC). Include a substrate blank containing 50 μ of assay buffer and 50 μ of 200 μM substrate. Then read at excitation and emission wavelengths of 380 nm and 460 nm, respectively, in kinetic mode for 5 minutes. The result was shown and it was obvious that PAI2 significantly decreased uPA activity. The inhibition IC50 was <300 nM.

Molecular Mass	47.9kDa
Endotoxin	<1.0EU per 1μg (determined by the LAL method).
Purity	> 95% as determined by SDS-PAGE.
Storage	Avoid repeated freeze/thaw cycles. Store at -20°C for 12 months. Aliquot and store at -80°C for 12 months.
Reconstitution	Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

GENE INFORMATION

Gene Name	Serpina2 serine (or cysteine) peptidase inhibitor, clade B, member 2 [<i>Mus musculus</i>]
Official Symbol	SERPINB2
Synonyms	SERPINB2; serine (or cysteine) peptidase inhibitor, clade B, member 2; plasminogen activator inhibitor 2, macrophage; serpin B2; plasminogen activator inhibitor, type II; serine (or cysteine) proteinase inhibitor, clade B, member 2; serine (or cysteine) proteinase inhibitor, clade B (ovalbumin), member 2; PAI-2; Planh2; ovalbumin;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID 18788


mRNA Refseq NM_001174170

Protein Refseq NP_001167641

Activity



SDS-PAGE



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA