

Recombinant Human XCL1 protein, His/Fc-tagged

Cat. No. XCL1-06H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human XCL1(Val22-Gly114) fused with His/Fc tag at C-terminal was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	Val22-Gly114
Description	Lymphotactin, also known as ATAC, C motif chemokine 1, Cytokine SCM-1, Lymphotaxin, SCM-1-alpha, Small-inducible cytokine C1, XC chemokine ligand 1, LTN, SCYC1 and XCL1, is a secreted protein which belongs to the intercrine gamma family. It is also a member of the XC chemokine family. XCL1 is found in spleen with highest level, lower in peripheral leukocytes and very low levels in lung, colon and small intestine. XCL1 plays a role in inflammatory and immunological responses, inducing leukocyte migration and activation. XCL1 induces chemotactic function by binding to a chemokine receptor called XCR1. XCL1 is closely related to another chemokine called XCL2.
Form	Lyophilized from a 0.2 µm filtered solution of PBS
AA Sequence	VGSEVSDKRTCVSLLTQRLPVSRIKTYTITEGSLRAVIFITKRGLKVCADPQATWVRD VVRSMRDR KSNTRNNMIQTKPTGTQQSTNTAVTLTGVDIEGRMDPEPKSCDKTHTC PPCPAPELLGGPSVFLF PPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPREEQYNSTYRVVSVLTV LHQDWLNGKEYKCKVSNKALPAPIEKTISKA

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

KGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFY PSDIAVEWESNGQPENNYKTT
PPVLDSGDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQ KSLSLSPGKHH
HHHH

Endotoxin Less than 0.1 ng/g (1 IEU/g).

Purity Greater than 95% as determined by SEC-HPLC and reducing SDS-PAGE.

Storage Lyophilized protein should be stored at < -20 centigrade, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7 centigrade for 2-7 days. Aliquots of reconstituted samples are stable at < -20 centigrade for 3 months.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in 1X PBS. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Shipping The product is shipped at ambient temperature.

GENE INFORMATION

Gene Name XCL1 chemokine (C motif) ligand 1 [Homo sapiens]

Official Symbol XCL1

Synonyms XCL1; chemokine (C motif) ligand 1; LTN, SCYC1, small inducible cytokine subfamily C, member 1 (lymphotactin); lymphotactin; ATAC; LPTN; SCM 1; SCM 1a; SCM-1-alpha; lymphotaxin; cytokine SCM-1; c motif chemokine 1; XC chemokine ligand 1; single cysteine motif 1a; small-inducible cytokine C1; small inducible cytokine subfamily C, member 1 (lymphotactin); LTN; SCM1; SCM-1; SCM1A; SCYC1; SCM-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	1a;
Gene ID	6375
mRNA Refseq	NM_002995
Protein Refseq	NP_002986
MIM	600250
UniProt ID	P47992
Chromosome Location	1q24.2
Pathway	Chemokine receptors bind chemokines, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Class A/1 (Rhodopsin-like receptors), organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; G alpha (q) signalling events, organism-specific biosystem;
Function	chemokine activity; chemokine receptor binding; chemokine receptor binding; protein homodimerization activity;