

Native Human CST3

Cat. No. CST3-26152TH **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Full length native protein isolated from urine (purified) (Human).
Species	Human
Description	The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. The type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of human fluids and secretions, where they appear to provide protective functions. The cystatin locus on chromosome 20 contains the majority of the type 2 cystatin genes and pseudogenes. This gene is located in the cystatin locus and encodes the most abundant extracellular inhibitor of cysteine proteases, which is found in high concentrations in biological fluids and is expressed in virtually all organs of the body. A mutation in this gene has been associated with amyloid angiopathy. Expression of this protein in vascular wall smooth muscle cells is severely reduced in both atherosclerotic and aneurysmal aortic lesions, establishing its role in vascular disease.
Tissue specificity	Expressed in submandibular and sublingual saliva but not in parotid saliva (at protein level). Expressed in various body fluids, such as the cerebrospinal fluid and plasma. Expressed in highest levels in the epididymis, vas deferens, brain, thymus, and ov
Form	Lyophilised: Use 100ul of a phosphate buffer, pH>7 containing 0.15 M NaCl to

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

reconstitute to 1mg/ml. Single homogenous batch, lyophilised from 0.02 M NH₄HCO₃ may contain traces of buffer salts.

Purity >95% by SDS-PAGE

Storage Store at +4°C.

Sequence Similarities Belongs to the cystatin family.

Full Length Full L.

GENE INFORMATION

Gene Name CST3 cystatin C [Homo sapiens]

Official Symbol CST3

Synonyms CST3; cystatin C; cystatin C (amyloid angiopathy and cerebral hemorrhage); cystatin-C;

Gene ID 1471

mRNA Refseq NM_000099

Protein Refseq NP_000090

MIM 604312

Uniprot ID P01034

Chromosome 20p11.2

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Location	
Pathway	Amyloids, organism-specific biosystem; Salivary secretion, organism-specific biosystem; Salivary secretion, conserved biosystem;
Function	beta-amyloid binding; cysteine-type endopeptidase inhibitor activity; peptidase inhibitor activity; protease binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA