

Recombinant Human DEFB4A protein

Cat. No. DEFB4A-27476TH **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human DEFB4A protein was expressed in Escherichia coli.
Species	Human
Source	E.coli
ProteinLength	41
Description	Defensins (alpha and beta) are cationic peptides with antimicrobial activity against Gram-negative and Gram-positive bacteria, fungi, and enveloped viruses. They are 2-6 kDa proteins and take important roles in innate immune system. On the basis of their size and pattern of disulfide bonding, mammalian defensins are classified into alpha, beta and theta categories. β-Defensins contain a six-cysteine motif that forms three intra-molecular disulfide bonds. Four human β-defensins have been identified and they are expressed on some leukocytes and at epithelial surfaces. Because β-defensins is cationic peptides, they can therefore interact with the membrane of invading microbes, which are negative due to lipopolysaccharides (LPS) and lipoteichoic acid (LTA) found in the cell membrane. Especially, they have higher affinity to the binding site compared to Ca^{2+} and Mg^{2+} ions. Furthermore, they can affect the stability of the membrane.
Form	Lyophilized from a 0.2 μm filtered concentrated solution in 20 mM PB, pH 7.4, 130 mM NaCl.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Bio-activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using immature human dendritic cells is in a concentration range of 10-100 ng/ml.
Molecular Mass	Approximately 4.3 kDa, a single non-glycosylated polypeptide chain containing 41 amino acids.
AA Sequence	GIGDPVTCLKSGAICHPVFCPRRYKQIGTCGLPGTKCCKKP
Endotoxin	Less than 1 EU/g of rHuBD-2 as determined by LAL method.
Purity	>98% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	DEFB4A
Official Symbol	DEFB4A
Synonyms	DEFB4A; defensin, beta 4A; DEFB2, DEFB4, DEFB102, defensin, beta 2 , defensin,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

beta 4; beta-defensin 4A; DEFB 2; HBD 2; SAP1; defensin, beta 2; defensin, beta 4; skin-antimicrobial peptide 1; BD-2; DEFB2; DEFB4; HBD-2; DEFB-2; DEFB102;

Gene ID 1673


mRNA Refseq NM_004942

Protein Refseq NP_004933

MIM 602215

UniProt ID O15263

**SDS-PAGE of
DEFB4A-27476TH**



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA