

Recombinant Human FAS protein

Cat. No. FAS-26332TH **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human FAS protein was expressed in Escherichia coli.
Species	Human
Source	E.coli
ProteinLength	157
Description	Fas and Fas Ligand (FasL) belong to the TNF superfamily and are type I and type II transmembrane proteins, respectively. Binding of FasL to Fas triggers apoptosis in Fas-bearing cells. The mechanism of apoptosis involves recruitment of pro-caspase 8 through an adaptor molecule called FADD followed by processing of the pro-enzyme to active forms. These active caspases then cleave various cellular substrates leading to the eventual cell death. sFasR is capable of inhibiting FasL-induced apoptosis by acting as a decoy receptor that serves as a sink for FasL.
Form	Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by its ability to inhibit the cytotoxicity of Jurkat cells is between 10-15 µg/ml in the presence of 2 ng/ml of rHuFas Ligand.
Molecular Mass	Approximately 17.6 kDa, a single non-glycosylated polypeptide chain containing 157 amino acids.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

AA Sequence	RLSSKSVNAQVTDINSKGLELRKTVTTVETQNLEGLHHDGQFCHKPCPPGERKARD CTVNGDEPDCVPCQEGKEYTDKAHFSSKCRRRCRLCDEGHGLEVEINCTRTQNTKC RCKPNFFCNSTVCEHCDPCTKCEHGIIECTLTSTNTKCKEEGSRSN
Endotoxin	Less than 1 EU/μg of rHusFasR/TNFRSF6 as determined by LAL method.
Purity	>95% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	FAS
Official Symbol	FAS
Synonyms	FAS; Fas (TNF receptor superfamily, member 6); APT1, FAS1, TNFRSF6, tumor necrosis factor receptor superfamily, member 6; tumor necrosis factor receptor superfamily member 6; APO 1; CD95; Fas AMA; FAS 827dupA; CD95 antigen; FASLG receptor; apoptosis antigen 1; Delta Fas/APO-1/CD95; APO-1 cell surface antigen; apoptosis-mediating surface antigen FAS; tumor necrosis factor receptor superfamily, member 6; APT1; FAS1; APO-1; FASTM; ALPS1A; TNFRSF6;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID	355
mRNA Refseq	NM_000043
Protein Refseq	NP_000034
MIM	134637
UniProt ID	P25445

**SDS-PAGE of FAS-
26332TH**

