

Recombinant Human Fibrillarin Protein, His-tagged

Cat. No. FBL-84H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human Fibrillarin protein, fused with N-terminal His tag, was expressed in E. coli.
Species	Human
Source	E.coli
ProteinLength	1-321
Description	Human recombinant fibrillarin (residues 2-321; Genbank Accession # NM_001436; MW= 35.1 kDa) expressed in E. coli with an N-terminal His-tag. Fibrillarin is a component of nucleolar ribonucleoprotein complexes including snoRNPs (small nucleolar ribonucleoproteins) and Cajal bodies. As part of the C/D snoRNPs, fibrillarin functions as the catalytic subunit in the 2'-O-methylation of ribosomal RNAs ¹ . In addition, fibrillarin has recently been shown to catalyze methylation of histone H2A glutamine-104, a modification restricted to the nucleolus and linked to RNA Pol I transcriptional activity on rDNA ³ . A distinct protein subcomplex comprising fibrillarin, SF2A-p32 and the arginine methyltransferases PRMT1 and PRMT5 has been described ⁴ and fibrillarin's N-terminal glycine and arginine-rich (GAR) domain, including in the context of full-length fibrillarin ⁵ , is a substrate for various PRMTs.
Form	50 mM Tris HCl, pH 7.5, 500 mM NaCl, 1 mM TCEP, 10 % glycerol
Molecular Mass	34.6 kDa

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Purity	>85% by SDS-PAGE
Notes	This product is NOT intended for therapeutic or diagnostic use in animals or in humans.
Storage	Store at -70 centigrade. Thaw quickly and store on ice before use. The remaining, unused, undiluted protein should be snap frozen, for example in a dry/ice ethanol bath or liquid nitrogen. Minimize freeze/thaws if possible, but very low volume aliquots (<5 μ l) or storage of diluted protein is not recommended.


GENE INFORMATION

Gene Name	FBL fibrillarin [Homo sapiens]
Official Symbol	FBL
Synonyms	FBL; fibrillarin; FIB; FLRN; RNU3IP1;
Gene ID	2091
mRNA Refseq	NM_001436
Protein Refseq	NP_001427
MIM	134795
UniProt ID	P22087


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



Coomassie bluestained SDS-PAGE (4-12% acrylamide) of 5 μ g of Fibrillarin (His). MW markers (left) are, from top, 220, 160, 120, 100, 90, 80, 70, 60, 50, 40, 30, 25, 20, 15, & 10 kDa.



Time courses of PRMT5/MEP50 Complex and PRMT5 (alone) methyltransferase reactions. Methylation determined as TCA-precipitable counts in a scintillation/filter plate assay. PRMT5/MEP50 Complex and PRMT5, at 30 nM and 15 nM respectively, were assayed with 0.4 μ M Fibrillarin (His) and 1 μ M [3H]-SAM.