

Recombinant Human IDH1(R132C) Protein, His-tagged

Cat. No. IDH1-111H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human IDH1(R132C) protein, fused with C-terminal His tag, was expressed in E.coli.
Species	Human
Source	E.coli
ProteinLength	1-414
Form	50 mM HEPES, pH 7.5, 200 mM NaCl, 3 mM DTT, 10% glycerol
Molecular Mass	47.7 kDa
Storage	Store at -80°C. Thaw quickly and store on ice before use. Avoid repeated freezing and thawing cycles.

GENE INFORMATION

Gene Name	IDH1 isocitrate dehydrogenase 1 (NADP+), soluble [Homo sapiens]
Official Symbol	IDH1
Synonyms	IDH1; isocitrate dehydrogenase 1 (NADP+), soluble; isocitrate dehydrogenase [NADP] cytoplasmic; NADP(+)-specific ICDH; oxalosuccinate decarboxylase; NADP-dependent isocitrate dehydrogenase, cytosolic; NADP-dependent isocitrate

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	dehydrogenase, peroxisomal; IDH; IDP; IDCD; IDPC; PICD;
Gene ID	3417
mRNA Refseq	NM_005896
Protein Refseq	NP_005887
MIM	147700
UniProt ID	O75874
Chromosome Location	2q32-qter
Pathway	Citrate cycle (TCA cycle), organism-specific biosystem; Citrate cycle (TCA cycle), conserved biosystem; Citrate cycle, first carbon oxidation, oxaloacetate =>2-oxoglutarate, organism-specific biosystem; Citrate cycle, first carbon oxidation, oxaloacetate => 2-oxoglutarate, conserved biosystem; Glutathione metabolism, organism-specific biosystem;
Function	NAD binding; isocitrate dehydrogenase (NADP+) activity; isocitrate dehydrogenase (NADP+) activity; magnesium ion binding; oxidoreductase activity; oxidoreductase activity, acting on the CH-OH group of donors, NAD or NADP as acceptor; protein homodimerization activity;