

Recombinant Human IDH2(R140Q) Protein, His-tagged

Cat. No. IDH2-116H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human IDH2(R140Q) protein, fused with C-terminal His tag, was expressed in E.coli.
Species	Human
Source	E.coli
ProteinLength	40-452
Form	50 mM HEPES, pH 7.5, 200 mM NaCl, 3 mM DTT, 10% glycerol
Molecular Mass	47.8 kDa
Purity	>90% by SDS-PAGE
Storage	Store at -80°C. Thaw quickly and store on ice before use. Avoid repeated freezing and thawing cycles.

GENE INFORMATION

Gene Name	IDH2 isocitrate dehydrogenase 2 (NADP+), mitochondrial [Homo sapiens]
Official Symbol	IDH2
Synonyms	IDH2; isocitrate dehydrogenase 2 (NADP+), mitochondrial; isocitrate dehydrogenase

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	[NADP], mitochondrial; NADP(+)-specific ICDH; oxalosuccinate decarboxylase; IDH; IDP; IDHM; IDPM; ICD-M; D2HGA2; mNADP-IDH;
Gene ID	3418
mRNA Refseq	NM_002168
Protein Refseq	NP_002159
MIM	147650
UniProt ID	P48735
Chromosome Location	15q21-qter
Pathway	Citrate cycle (TCA cycle), organism-specific biosystem; Citrate cycle (TCA cycle), conserved biosystem; Citrate cycle, first carbon oxidation, oxaloacetate =>2-oxoglutarate, organism-specific biosystem; Citrate cycle, first carbon oxidation, oxaloacetate => 2-oxoglutarate, conserved biosystem; Citric acid cycle (TCA cycle), organism-specific biosystem;
Function	NAD binding; isocitrate dehydrogenase (NADP+) activity; isocitrate dehydrogenase (NADP+) activity; magnesium ion binding; oxidoreductase activity; oxidoreductase activity, acting on the CH-OH group of donors, NAD or NADP as acceptor;