

Recombinant Human IL12A protein

Cat. No. IL12A-26973TH Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human IL12A protein was expressed in CHO.
Species	Human
Source	CHO
ProteinLength	525
Description	Interleukin-12 (IL-12), also known as NKSF or CLMF, is a pleiotropic cytokine originally identified in the medium of activated human B lymphoblastoid cell lines. The p40 subunit of IL-12 has been shown to have extensive amino acid sequence homology to the extracellular domain of the human IL-6 receptor while the p35 subunit shows distant but significant sequence similarity to IL-6, G-CSF, and chicken MGF. These observations have led to the suggestion that IL-12 might have evolved from a cytokine/soluble receptor complex. Human and murine IL-12 share 70 % and 60 % amino acid sequence homology in their p40 and p35 subunits, respectively. IL-12 apparently shows species specificity with human IL-12 reportedly showing minimal activity in the murine system.
Form	Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by a cell proliferation assay using PHA-activated human T lymphoblasts is less than 0.05 ng/ml, corresponding to a specific activity of $> 2.0 \times 10^7$ IU/mg.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	Approximately 75 kDa, consisting of a 306 amino acid residue p40 subunit and a 197 amino acid residue p35 subunit.
AA Sequence	p35Subunit:RNLPVATPDPMFCLHHSQNLLRAVSNMLQKARQTLEFYPTCTSEEID HEDITKDKTSTVEACLPLELTKNESCLNSRETSFITNGSCLASRKTSFMMALCLSSIYE DLKMYQVEFKTMNAKLLMDPKRQIFLDQNMLAVIDELMQALNFNSETVPQKSSLEEP DFYKTKIKLCILLHAFRIRAVTIDRVMSYLNASp40Subunit:IWELKKDVYVVELDWYPD APGEMVVLTCDTPEEDGITWTLTQSSEVLGSGKTLTIQVKEFGDAGQYTCHKGGEV LSHSLLLLHKKEDGIWSTDILKDQKEPKNKTFLRCEAKNYSGRFTCWWLTTISTDLTF SVKSSRGSSDPQGVTCGAATLSAERVGRDGNKEYEYSVEQCEDSACPAAEESLPIEV MVDVHKLKYENYTSSFFIRDIKPDPPKNLQLKPLKNSRQVEVSWEYPDTWSTPHS YFSLTFCVQVQGKSKREKKDRVFTDKTSATVICRKNASISVRAQDRYYSSSWSEWA SVPCS
Endotoxin	Less than 1 EU/g of rHuIL-12 as determined by LAL method.
Purity	>95% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene Name	IL12A
Official Symbol	IL12A
Synonyms	IL12A; interleukin 12A (natural killer cell stimulatory factor 1, cytotoxic lymphocyte maturation factor 1, p35); NKSF1; interleukin-12 subunit alpha; CLMF; cytotoxic lymphocyte maturation factor 1; p35; IL 12; subunit p35; IL 12A; IL35 subunit; interleukin 12; interleukin 12 alpha chain; natural killer cell stimulatory factor 1; 35 kD subunit; NF cell stimulatory factor chain 1; NFSK; CLMF p35; IL-12 subunit p35; IL-12, subunit p35; interleukin 12, p35; interleukin-12 alpha chain; NK cell stimulatory factor chain 1; cytotoxic lymphocyte maturation factor 1, p35; cytotoxic lymphocyte maturation factor 35 kDa subunit; natural killer cell stimulatory factor 1, 35 kD subunit; P35; IL-12A;
Gene ID	3592
mRNA Refseq	NM_000882
Protein Refseq	NP_000873
MIM	161560
UniProt ID	P29459

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA