Fluro IndoBlu cell proliferation Detection Kit

Cat. No. Kit-0204 **Lot. No.** (See product label)

SPECIFICATION

Product Overview	
Description	Measurement of cell proliferation and viability is frequently used in clinical and experimental immunology as means of assessing cell activation in response to diseases, infections and environmental stimulations. Fluoro IndoBlu assay can be used for the measurement of cell proliferation in response to antigens, cytokines, growth factors, and mitogens. It can also be used for the analysis of cytotoxic effects of anticancer drugs, drug resistance, cytotoxic pharmaceutical compounds, and other toxic agents.Staurosporine induced Jurkat cell viability assay. Titrated doses of staurosporine were added into seeded Jurkat (10,000 cells/well) cells in 96 well black opaque tissue culture plates and incubated at 37°C, 10% CO2 for 24h. Fluoro IndoBluTM (1/10 volume) was added into the cultured cells and incubated for 3h and fluorescence was detected at Ex: 530nm and Em: 590nm.PBMC proliferation in response to mitogen Concanavalin A (Con A) stimulation measured by Fluoro IndoBluTM. PBMC at 15,000 cells were cultured for 3 days in the presence of titrated amounts of Con-A and tested for proliferation using the Fluoro IndoBluTM assay. Fluoro IndoBluTM (1/10 volume) was added to the cultured cells and incubated 37°C for 2h. Fluorescence was detected at Ex: 530nm and Em: 590nm.
Applications	Fluorescence plate reader
Usage	1. This kit is only for usage on cell lines or primary cell cultures.2. For research use only. Not for use in diagnostic procedures.3. Gloves, protective clothing and eyewear should be worn and safe laboratory practices followed.

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA

Storage	1. Short term: Store the kit at 2-8°C for (1 -2 weeks), keep it away from light.2. Long term: Store the kit at -20°C.
Kit Components	1. IndoBlu Reagent: 2-8°C;2. Optional: 96-Well Clear Bottomed Black Plate for Fluorescent Readout
Features & Benefits	1. Highly effective and stable assay for cell proliferation.2. No need to wash out media from cell samples, just add the reagent directly to your experimental samples.3. Works for both cell viability and proliferation assay.4. Plate can be read in 30 minutes to 3 hours.

Tel: 1-631-559-9269 1-516-512-3133
Email: info@creative-biomart.com Fax: 1-631-938-8127
45-1 Ramsey Road, Shirley, NY 11967, USA