

Di-Methyl Histone H4K20 Quantification Kit (Colorimetric)

Cat. No. Kit-0289 **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Di-Methyl Histone H4K20 Quantification Kit (Colorimetric) is used for measuring di-methylation of histone H4K20.

Description

Epigenetic activation or inactivation of genes play a critical role in many important human diseases, especially in cancer. A major mechanism for epigenetic inactivation of the genes is methylation of CpG islands in genome DNA caused by DNA methyltransferases. Histone methyltransferases (HMTs) control or regulate DNA methylation through chromatin-dependent transcriptional repression or activation. HMTs transfer 1-3 methyl groups from S-adenosyl-L-methionine to the lysine and arginine residues of histone proteins. PR-SET7, SET9, SUV4.20h, and ASH1 are histone methyltransferases that catalyze methylation of histone H4 at lysine 20 (H4K20) in mammalian cells. Di-methylation of H4K20 has been described as another repressive chromatin domain and is involved in DNA damage response. The H4K20 di-methylation can also be changed by inhibition or activation of HMTs, making quantitative detection of di-methyl histone H4K20 a useful tool for better understanding epigenetic regulation of gene activation/repression. This information is also useful for developing HMT-targeted drugs. Di-Methyl Histone H4K20 Quantification Kit (Colorimetric) provides a tool for measuring di-methylation of histone H4K20.

Applications

Di-Methyl Histone H4K20 Quantification Kit (Colorimetric) is suitable for specifically measuring histone H4K20 di-methylation using a variety of mammalian cells (human, mouse, etc.) including fresh and frozen tissues, cultured adherent and suspension cells.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Usage	For research use only (RUO)
Storage	Upon receipt, store standard control at –20°C. Store all other components at 4°C away from light. The components of the kit are stable for up to 6 months from date of shipment when stored properly. Note: Check if buffers C1 and C2 contain salt precipitates before using. If so, warm (at room temperature or 37°C) and shake the buffers until the salts are re-dissolved.
Kit Components	C1 (10X wash buffer) 10 mL C2 (antibody buffer) 6 mL C3 (detecting antibody, 1 mg/mL)* 5 µL C4 (color developer) 5 mL C5 (stop solution) 3 mL Standard control (100 µg/mL)* 10 µL Signal report solution* 5 µL Signal enhancer* 120 µL 8 well sample strips (with frame) 48 well standard control strips 2* For maximum recovery of the products, centrifuge the original vial prior to opening the cap.
Detection method	Colorimetric
Compatible Sample Types	Histone Extract
Features & Benefits	Quick and efficient procedure, which can be finished within 2.5 hours. Innovative colorimetric assay without the need for radioactivity, electrophoresis, or chromatography. Specifically captures di-methylated H4K20 with the detection limit as low as 5 ng/well and detection range from 20 ng-2 µg/well of histone extracts. The control is conveniently included for quantification of di-methylated H4K20. Strip microplate format makes the assay flexible: manual or high throughput. Simple, reliable, and consistent assay conditions.