

HDAC Activity/Inhibition Colorimetric Assay Kit

Cat. No. Kit-0424 **Lot. No.** (See product label)

SPECIFICATION

Product Overview	HDAC Activity/Inhibition Assay Kit (Colorimetric) is use for measuring HDAC activity/inhibition.
Description	<p>Histone deacetylases (HDACs) play a critical role in transcriptional repression of the gene expression in eukaryotic cells through catalyzing the hydrolytic removal of acetyl groups from histone lysine residues. HDACs are tightly involved in cell cycle regulation, cell proliferation, and in development of human cancer. HDAC inhibition displays significant effects on apoptosis, cell cycle arrest, and differentiation in cancer cells. HDAC inhibitors are currently being developed as potential anticancer agents. There are several methods used for measuring HDAC activity/ inhibition. However most of these methods available so far are time consuming, laborious, produce radioactive waste, or cannot measure precise HDAC activity and inhibitory effects of inhibitors.</p>
Applications	For measuring HDAC activity/inhibition from a broad range of species including mammalian cells/tissues, plants, and bacteria.
Usage	For research use only (RUO)
Storage	Upon receipt, store H3, H4, H5 and H7 at –20°C away from light. Store H6, H8 and 8 well assay strips at 4°C away from light. Store all other components at room temperature. The components of the kit should be stable for 6 months when stored properly.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Kit Components

H1 (10X wash buffer) 28 ml H2 (HDAC assay buffer) 3 ml H3 (biotinylated HDAC substrate)* 100 μ l H4 (HDAC inhibitor, 0.5 mM)* 100 μ l H5 (HDAC standard, 20 μ g/ml)* 50 μ l H6 (capture antibody 1000 μ g/ml)* 50 μ l H7 (detection antibody 200 μ g/ml)* 20 μ l H8 (developing solution) 12 ml H9 (stop solution) 6 ml 8 well assay strip (with frame) 12*

For maximum recovery of the products, centrifuge the original vial after thawing prior to opening the cap.

Features & Benefits

Fast procedure, which can be finished within 3 hours. Innovative colorimetric assay without need for radioactivity, extraction, and chromatography. Direct measurement of HDAC activity and inhibition with no use of lysyl endopeptidase, thereby avoiding the false inhibitory effect on HDACs and allowing more accurate measurement. Strip microplate format makes the assay flexible: manual or high throughput analysis. Simple, reliable, and consistent assay conditions.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA