

LSD1 Demethylase Activity/Inhibition Fluorometric Assay Kit

Cat. No. Kit-0524 **Lot. No.** (See product label)

SPECIFICATION

Product Overview

LSD1 Demethylase Activity/Inhibition Assay Kit (Fluorometric) is used for an easy and fast fluorometric measurement of LSD1 activity or inhibition.

Description

Lysine histone methylation is one of the most robust epigenetic marks and is essential for the regulation of multiple cellular processes. The methylation of H3-K4 seems to be of particular significance, as it is associated with active regions of the genome. H3-K4 methylation was considered irreversible until the identification of a large number of histone demethylases indicated that demethylation events play an important role in histone modification dynamics. So far at least 2 classes of H3-K4 specific histone demethylase, LSD1 (BHC110, KDM1) and JARIDs have been identified. LSD1 can remove di- and mono-methylation from H3-K4 by using an amine oxidase reaction. LSD1 is associated with complexes that function as both transcriptional inactivators and activators. It demethylates mono-/di-methyl H3-K4 when associated with the Co-REST complex at neuronal genes, or mono-/di-methyl H3-K9 when associated with the androgen receptor.

Applications

The LSD1 Demethylase Activity/Inhibition Assay Kit (Fluorometric) is suitable for measuring LSD1 activity/inhibition using nuclear extracts or purified enzymes from a broad range of species such as mammals, plants, fungi, and bacteria, in a variety of forms including cultured cells and fresh tissues. Nuclear extracts can be prepared by using your own successful method. Nuclear extracts can be used immediately or stored at -80°C for future use. Purified enzymes can be active LSD1 from recombinant proteins or isolated from cell/tissues.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Usage	For research use only (RUO)
Storage	Upon receipt: (1) Store LF3, LF4, LF6 and LF8 at –20°C away from light; (2) Store LF1, LF5, LF7, LF9, and 8-Well Assay Strips at 4°C away from light; (3) Store remaining components (LF2, LF10, and Adhesive Covering Film) at room temperature away from light. All components of the kit are stable for 6 months from the date of shipment, when stored properly. Note: (1) Check if LF1 (10X Wash Buffer) contains salt precipitates before use. If so, warm (at room temperature or 37°C) and shake the buffer until the salts are re-dissolved.
Kit Components	LF1 (10X Wash Buffer) 14 mL LF2 (LSD1 Assay Buffer) 4 mL LF3 (LSD1 Substrate, 50 µg/mL)* 60 µL LF4 (LSD1 Assay Standard, 50 µg/mL)* 10 µL LF5 (Capture Antibody, 1000 µg/mL*) 5 µL LF6 (Detection Antibody, 400 µg/mL)* 6 µL LF7 (LSD1 Inhibitor Tranylcypromine, 1 mM)* 20 µL LF8 (Fluoro Developer) 10 µL LF9 (Fluoro Enhancer) 10 µL LF10 (Fluoro Diluter) 4 mL 8-Well Assay Strips (With Frame) 6 strips Adhesive Covering Film 1 slice* Spin the solution down to the bottom prior to use.
Detection method	Fluorometric
Features & Benefits	Strip-well microplate format makes the assay flexible and quick: manual or high throughput analysis that can be completed within 3 hours. Enhanced kit composition enables background signals to be extremely low, which allows the assay to be more accurate, sensitive, reliable, and consistent. Innovative fluorometric assay directly measures LSD1 activity by a straightforward detection of LSD1-converted demethylated product, rather than by-products. Thus it eliminates assay interferences caused by thiol-containing chemicals such as DTT, GSH, and 2-mercaptoethanol. Both cell/tissue extracts and purified LSD1 can be used, which allows for the detection of inhibitory effects of LSD1 inhibitor in vivo and in vitro. Novel assay principle allows high sensitivity to be achieved. The activity can be detected from as low as 2 ng of purified LSD1 enzyme, which is about 50 fold higher than that

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

obtained by H₂O₂/formaldehyde release-based LSD1 assays. Demethylated H3-K4 standard is included, which allows the specific activity of LSD1 to be quantified.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA