

Cell-Free Protein Expression Kit

Cat. No. Kit-0883 **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Cell-Free Protein Expression Kit is designed for in vitro transcription and translation of target DNA to protein in a single reaction. This product contains an optimized E.coli extract containing T7 RNA polymerase for transcription and ribosome for translation. An optimized Master mix provides all other required components, including amino acids, rNTPs, and appropriate salts to express high-level production of recombinant proteins. This product contains reagents for expression of protein directly from a variety of DNA templates, including T7 promoter and terminator as well as RBS (ribosomal binding site). This product can be used to express a wide range of target proteins and has the capacity for a yield of more than 300 µg/mL of protein within 3 hours
Applications	This product can be used to express a wide range of target proteins and has the capacity for a yield of more than 300 µg/mL of protein within 3 hours.
Storage	Store kit at -20°C ~ -70°C.
Size	45 µL x 24 reactions
Kit Components	Master mix: 0.54 mL x 1 tube; E.coli extract: 0.3 mL x 1 tube; DEPC DW: 1 mL x 1 tube; Positive Control DNA: 1 tube.
Features & Benefits	High-speed: synthesize your target protein quickly (within 3 hours) and economically

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Easy-to-Use: includes all necessary components for transcription and translation

Flexible: synthesize proteins from various DNA templates.

synthesize proteins of toxic proteins

incorporate non-natural amino acids

use additives such as metal ions, detergents, cofactors, binding partners

Assay Protocol

1. Thaw components in the Cell-Free Protein Expression kit on ice.
2. After thawing, spin-down the the Master mix and E.coli extract, and then place the tube on ice bath.
3. Prepare the reaction mixture (one reaction).
 Negative: Master mix 21 μ ; E.coli extract 12 μ ; DEPC DW 12 μ ; Total 45 μ .
 Positive: Master mix 21 μ ; E.coli extract 12 μ ; DNA template 1 μ ; DEPC DW 11 μ ; Total 45 μ .
 Sample: Master mix 21 μ ; E.coli extract 12 μ ; DNA template X μ ; DEPC DW (12-X) μ ; Total 45 μ .
4. Mix reaction mixture by tapping the tube gently and spin down briefly to collect the mixture at the bottom of the tube.
5. Incubate the reaction mixture at 30°C in a water bath for 3 hours.
6. After completing incubation, spin down briefly to collect the mixture at the bottom of the tube.
7. Analyze the mixture using SDS-PAGE, western blotting or bioactivity assay.

Analysis

- After expression of proteins using Cell-Free Protein Expression Kit, you can the expression of target protein via SDSPAGE
1. Prepare loading samples.
 5 μ of expression sample + 10 μ of D.W. + 5 μ of 4x Loading dye (Final volume is 20 uL)
 2. Denature all samples at 95°C for 5 min.
 3. Load 5 μ of each samples on SDS-PAGE minigel.
 4. Run SDS-PAGE gel.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

5. Perform staining with coomassie blue R-250.

Ex) Result of positive control (28kDa)

-condition : 12% SDS-PAGE gel, at 50 V for 20 min→at 150 V for 50 min

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA