

NAD/NADH Assay Kit

Cat. No. Kit-0613 **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Detection and Quantification of NAD/NADH Content.
Description	Nicotinamide adenine dinucleotide (NAD ⁺) is a vital coenzyme found in all cells. As NAD ⁺ is involved in redox reactions, it is found in two forms in cells. NAD ⁺ is an oxidizing agent and becomes reduced to form NADH, which can be used as a reducing agent. As a result, it plays a key role in metabolism and other cellular processes. In organisms, NAD ⁺ can be synthesized de novo from tryptophan or aspartic acid. Because of the wide variety of functions that NAD ⁺ plays, it is a popular target for pharmaceuticals. NAD/NADH Assay Kit is based on a glucose dehydrogenase cycling reaction, in which the formed NADH reduces a formazan (MTT) reagent. The intensity of the reduced product color, measured at 570 nm, is proportionate to the NAD ⁺ /NADH concentration in the sample.
Applications	Functional Assay
Storage	Shipped and store at 4°C for 6 months.
Kit Components	Component-Volume-Storage 1. 96-Well Microplate; 2. Extract I: 30 ml x 1, 4°C; 3. Extract II: 30 ml x 1, 4°C; 4. Reagent I: 10 ml x 1, 4°C; 5. Reagent II: 1.5 ml x 2, 4°C; 6. Reagent III: Powder x 1, -20°C; 7. Reagent IV: Powder x 1, 4°C; 8. Reagent V: 1.2 ml x 3, 4°C; 9. Reagent VI: 30 ml x 1, 4°C; 10. Reagent VII: 25 ml x 2, 4°C; 11. Technical Manual: 1 Manual Note: Reagent III: add 3 ml distilled water to dissolve before use, mix, stored at 4°C. Reagent IV: add 3 ml distilled water to dissolve before use, mix, stored at 4°C.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Detection method	Colorimetric
Compatible Sample Types	Urine; Serum; Plasma; Other biological fluids; Tissue Extracts; Cell Lysate