

1,5-Anhydroglucitol Uptake Assay Kit (Cell-Based)

Cat. No. Kit-1004 Lot. No. (See product label)

SPECIFICATION

Product Overview

Description

1,5-Anhydroglucitol (1,5-AG) is the 1-deoxy form of D-glucose. Its level in blood plasma has been used in clinical research to investigate short-term glycemic control levels in diabetic patients. Cereals and animal source proteins constitute the main 1,5-AG source for humans. Most of 1,5-AG is absorbed by renal tubules, and its levels in human body are highly regulated and relatively consistent. Glucose can act as a competitive inhibitor of 1,5-AG in blood. Therefore, glucose levels are inversely proportional to 1,5-AG concentrations. The physiological function of 1,5-AG is not well understood. Recent studies showed 1,5-AG uptake could be associated with sodium-dependent glucose transporter. 1,5-Anhydroglucitol Uptake Assay Kit utilizes a proprietary fluorescent 1,5-AG analog, which can be taken up by cells. However, this 1,5-AG analog cannot be fully utilized in glycolytic processes and thus accumulates inside the cells. Fluorescence generated by this analog is proportional to the cellular 1,5-AG uptake. The assay can be used to monitor 1,5-AG uptake by using fluorescent microscopy and flow cytometry. Phloretin, a compound that can inhibit glucose transporter and 1,5-AG cell uptake, is included in the kit as a control. This non-radioactive assay kit is easy-to-use and allows qualitative and quantitative measurements of 1,5-AG uptake in cultured cells.

Applications

Measurement of 1,5-Anhydroglucitol uptake in response to insulin, growth factors, cytokines, mitogens and nutrients, etc.

Dual-staining of 1,5-AG transporters and 1,5-AG uptake

Analysis of 1,5-AG metabolism and cell signaling in various cell types

Screening of anti-diabetic compounds

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage	-20°C
Shipping	Gel Pack
Size	50 assays
Kit Components	Analysis Buffer; AGTracker; Reagent (100X); Phloretin (100X) (in DMSO); Phloretin (100X)
Detection method	FACS (488 nm excitation laser) and fluorescent microscope (excitation range 420 nm-495 nm)
Features & Benefits	Easy-to-use; Non-radioactive; Image and accurately measure 1,5-Anhydroglucitol uptake in cultured cells in response to insulin, growth factors etc.

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA