

TR-FRET TR alpha Coactivator Assay Kit, goat

Cat. No. Kit-0897 **Lot. No.** (See product label)

SPECIFICATION

Product Overview

The TR-FRET TR (thyroid receptor) alpha Coactivator Assay Kit provides a sensitive and robust method for highthroughput screening of potential TR alpha ligands as agonists of ligand-dependent coactivator recruitment. The kit uses a terbium (Tb)-labeled anti-GST antibody, a fluorescein-labeled coactivator peptide, and a human recombinant TR alpha ligand-binding domain (TR alpha-LBD) that is tagged with glutathione-S-transferase (GST) in a homogeneous, mix-and-read assay format.

Description To Assay:
When performing the TR-FRET TR alpha coactivator assay, TR alpha-LBD is added to ligand test compounds followed by addition of a mixture of the fluorescein-coactivator peptide and Tb anti-GST antibody. After an incubation period at room temperature, the 520 nm/495 nm TR-FRET ratio is calculated and used to determine the EC₅₀ from a dose response curve of the compound. Based on the biology of the TR alpha-coactivator peptide interaction, the ligand EC₅₀ is a composite value representing the amount of ligand required to bind to receptor, induce a conformational change, and recruit coactivator peptide.

Applications Co-Factor Interaction Assay

Storage Store in ultra-cold freezer (-68 to -85°C).

Shipping Dry Ice

Size 800 x 20 µ assays

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Kit Components	The TR-FRET TR alpha Coactivator Assay Kit contains TR alpha-LBD (GST) protein, fluorescently labeled SRC2-2 coactivator peptide, Tb-labeled anti-GST antibody, and buffers. Store components as indicated in the assay protocol (-80°C, -20°C, or +4°C).
Materials Required but Not Supplied	Microplate Reader
Detection method	Fluorescent

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA