

Cyclic Nucleotide Phosphodiesterase assay Kit

Cat. No. Kit-0667 **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Unique, non-radioactive, colorimetric assay. Used to screen inhibitors and modulators of cyclic nucleotide phosphodiesterase activity. HTS format permitting rapid assays of large numbers of samples.
Size	96 wells
Description	A BIOMOL GREEN Quantizyme Assay System. The Cyclic Nucleotide Phosphodiesterase (PDE) Assay Kit combines a special dual enzyme system with the BIOMOL GREEN Reagent for phosphate detection to create a unique, non-radioactive, colorimetric assay. This HTS-friendly, mix and read system may be used to screen inhibitors and modulators of cyclic nucleotide phosphodiesterase activity. 96-well microplate format permits rapid assays of large numbers of samples. The basis for the assay is the cleavage of cAMP or cGMP by a cyclic nucleotide phosphodiesterase. The 5'-nucleotide released is further cleaved into the nucleoside and phosphate by the enzyme 5'-nucleotidase. The phosphate released due to enzymatic cleavage is quantified using BIOMOL GREEN reagent in a modified Malachite Green assay. The kit includes Type I cyclic AMP phosphodiesterase (PDE) for validation purposes.
Applications	Colorimetric detection, HTS
Storage	-80°C
Kit Components	PDE Enzyme (from bovine brain) (5 x 4 U; lyophilized solid, 4 U per vial. 1 U = 1 nmol

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

3",5-cAMP to 5"-AMP per minute under the conditions of the linearity assay (30°C, pH 7.4, 200 μ M 3",5"-cAMP Storage: -70°C, avoid freeze/thaw cycles.5"-Nucleotidase (1ml; 5 kU/ μ in 10mM TRIS-hydrochloric acid, pH 7.4, 125mM sodium chloride, 10% glycerol, 1mM magnesium chloride. One U will release 1 pmol phosphate per minute from 200 μ M 5"-AMP, 30°C in.a reaction buffer of 10mM TRIS-hydrochloric acid, pH 7.4, 0.2mM magnesium chloride). Storage: -70°C3",5"-cAMP Substrate (2ml; 1mM in assay buffer (10mM TRIS-hydrochloric acid, pH 7.4). Storage: -20 or -70°C3",5"-cGMP Substrate (2ml; 1mM in assay buffer (10mM TRIS-hydrochloric acid, pH 7.4). Storage: -20 or -70°C

CPDE Assay Buffer (40ml (2 x 20ml); 10mM TRIS-hydrochloric acid, pH 7.4) Storage: -70°C

Biomol Green Reagent (20ml; liquid in screw-cap plastic bottle). Storage: 4°C, long-term at -70°C

5"-AMP Standard (1.0ml; 100 μ M in 10mM TRIS-hydrochloric acid, pH 7.4 buffer). Storage: -20 or -70°C

5"-GMP Standard (1.0ml; 100 μ M in 10mM TRIS-hydrochloric acid, pH 7.4 buffer). Storage: -20 or -70°C

Inhibitor (IBMX) (200 μ l 200 μ M in assay buffer (10mM TRIS-hydrochloric acid, pH 7.4)). Storage: -20 or -70°C

Desalting column and resin (1 column and 1 g of resin). Storage: room temperature, after rehydration store at 4°C

½ volume microplate. Storage: Room temperature