

Recombinant Human IDH1 protein, DDK-tagged

Cat. No. IDH1-5849H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Human IDH1 fused with DDK tag at C-terminal was expressed in Insect cells.
Species	Human
Source	Insect Cells
Form	50mM Tris-HCl pH8.0, 150mM NaCl, 10%glycerol. Store at -80C. Avoid repeated freeze-thaw cycles. Stable for at least 3 months from receipt of products under proper storage and handling conditions.
Molecular Mass	47 kDa
Purity	> 80% as determined by SDS-PAGE and Coomassie blue staining
Concentration	>50 ug/mL as determined by microplate BCA method

GENE INFORMATION

Gene Name	IDH1 isocitrate dehydrogenase 1 (NADP+), soluble [Homo sapiens]
Official Symbol	IDH1
Synonyms	IDH1; isocitrate dehydrogenase 1 (NADP+), soluble; isocitrate dehydrogenase [NADP] cytoplasmic; NADP(+)-specific ICDH; oxalosuccinate decarboxylase; NADP-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	dependent isocitrate dehydrogenase, cytosolic; NADP-dependent isocitrate dehydrogenase, peroxisomal; IDH; IDP; IDCD; IDPC; PICD;
Gene ID	3417
mRNA Refseq	NM_005896
Protein Refseq	NP_005887
MIM	147700
UniProt ID	O75874
Chromosome Location	2q32-qter
Pathway	Citrate cycle (TCA cycle), organism-specific biosystem; Citrate cycle (TCA cycle), conserved biosystem; Citrate cycle, first carbon oxidation, oxaloacetate => 2-oxoglutarate, organism-specific biosystem; Citrate cycle, first carbon oxidation, oxaloacetate => 2-oxoglutarate, conserved biosystem; Glutathione metabolism, organism-specific biosystem;
Function	NAD binding; isocitrate dehydrogenase (NADP+) activity; isocitrate dehydrogenase (NADP+) activity; magnesium ion binding; oxidoreductase activity; oxidoreductase activity, acting on the CH-OH group of donors, NAD or NADP as acceptor; protein homodimerization activity;