

Recombinant Human NCOA1 protein, His-tagged and Biotinylated

Cat. No. NCOA1-372H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	The His-tag recombinant protein is purified by affinity chromatography in combination with FPLC columns.
Species	Human
Source	E.coli
Description	Steroid receptor coactivator 1 (SRC1) is a transcriptional coactivator that mediates the activating functions of many of the nuclear hormone receptors. It is also known as NCoA1 and is a member of the SRC/p160 coactivator family. SRC1 has been shown to be over expressed in some cancers. SRC1 is a 160 kDa protein that contains several LXXLL motifs, which are involved in nuclear receptor interaction. The region 627-786 contains 3 LXXLL motifs that are involved in interaction with nuclear hormone receptors and has been previously used in assays detecting ligand-dependent receptor-cofactor interactions. This protein does not contain any tag and will be useful for assays where a native untagged protein is desired.
Form	The protein is in 20mM Tris-HCl pH7.9, 100mM NaCl, 0.2mM EDTA, 1mM DTT and 20% glycerol.
Molecular Mass	21.9 kDa
AA Sequence	SQTSHKLVQL LTTTAEQQLR HADIDTSCKD VLSCTGTSNS ASANSSGGSC PSSHSSLTERHKILHRLLEQ GSPSDITLTS VEPDKKDSAS TSVSVTGQVQ

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	GNSSIKLELD ASKKKESKDHQLRLYLLDKD EKDLRSTPNL SLDDVKVKVE KKEQMDPCNT
Purity	greater than 90% homogeneous based on SDS-PAGE analysis
Unit Definition	1 unit equals 1 nanogram of purified protein. 100 units are sufficient for a protein-protein interaction assay.
Applications	The biotinylated SCR1-RID (627-786) can be used for protein-protein interaction assays, particularly suitable for high throughput screening (HTS) assays.
Storage	Stored at -70 centigrade before use. Avoid repeated freeze thaw cycles.
Conjugation	Biotin

GENE INFORMATION

Gene Name	NCOA1 nuclear receptor coactivator 1 [Homo sapiens]
Official Symbol	NCOA1
Synonyms	NCOA1; nuclear receptor coactivator 1; bHLHe74; F SRC 1; KAT13A; NCoA 1; RIP160; SRC1; Hin-2 protein; PAX3/NCOA1 fusion protein; steroid receptor coactivator-1; renal carcinoma antigen NY-REN-52; class E basic helix-loop-helix protein 74; F-SRC-1; bHLHe42; MGC129719; MGC129720;
Gene ID	8648
mRNA Refseq	NM_003743
Protein Refseq	NP_003734

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

MIM	602691
UniProt ID	Q15788
Chromosome Location	2p23
Pathway	Adipogenesis, organism-specific biosystem; Androgen Receptor Signaling Pathway, organism-specific biosystem; BMAL1:CLOCK/NPAS2 Activates Circadian expression, organism-specific biosystem; Circadian Clock, organism-specific biosystem; Circadian Repression of expression by REV-ERBA, organism-specific biosystem
Function	DNA binding; androgen receptor binding; chromatin binding; enzyme binding; histone acetyltransferase activity; ligand-dependent nuclear receptor binding; ligand-dependent nuclear receptor transcription coactivator activity; nuclear hormone receptor binding; protein N-terminus binding; protein binding; signal transducer activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA