

Recombinant Human NAA20 protein, T7-tagged

Cat. No. NAA20-171H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human NAA20 (178 aa, Isoform-A) fused with T7 Tag at N-terminal was expressed in E. coli.
Species	Human
Source	E.coli
ProteinLength	178 a.a.
Form	1.0 mg/ml, sterile-filtered, in 20 mM pH 8.0 Tris-HCl Buffer, with proprietary formulation of NaCl, KCl, EDTA, Sucrose and DTT.
AA Sequence	MASMTGGQQMGRGEFMTTLRAFTCDDLFRFNNINLDPLTETYGIPFYLQYLAHWPE YFIVAEAPGGELMGYIMGK AEGSVAREEWHGHVTALSVAPEFRRLGLAAKLMELLE EISERKGGFFVDLFVRVSNQVAVNMYKQLGYSVYRTVI EYYSASNGEPDEDAYDMR KALSRDTEKKSIPLPHPVRPEDIE
Purity	>90% by SDS-PAGE
Applications	1. May be used for in vitro NatB mediated cell cycle regulation study with intracellular delivery of this protein.2. As soluble / native protein, may be used as enzymatic substrate protein for kinase and ubiquitin assay development.3. May be used for mapping NAA20 protein-protein interaction.4. Potential diagnostic biomarker for cancer.5. May be used as antigen for specific antibody development.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage

Keep at -80 centigrade for long term storage. Product is stable at 4 centigrade for at least 7 days.

GENE INFORMATION
Gene Name

NAA20 N(alpha)-acetyltransferase 20, NatB catalytic subunit [Homo sapiens]

Official Symbol

NAA20

Synonyms

NAA20; N(alpha)-acetyltransferase 20, NatB catalytic subunit; N acetyltransferase 5 , N acetyltransferase 5 (ARD1 homolog, S. cerevisiae) , N acetyltransferase 5 (GCN5 related, putative) , N acetyltransferase 5, ARD1 subunit (arrest defective 1, S. cerevisiae, homolog) , NAT5; N-alpha-acetyltransferase 20; dJ1002M8.1; N acetyltransferase 3 homolog (S. cerevisiae); NAT3; natB catalytic subunit; natB complex subunit NAT5; N-acetyltransferase 3 homolog; N-acetyltransferase 5 (GCN5-related, putative); N-terminal acetyltransferase complex ARD1 subunit; N-acetyltransferase 5 (ARD1 homolog, S. cerevisiae); N-alpha-acetyltransferase 20, NatB catalytic subunit; N-terminal acetyltransferase B complex catalytic subunit NAT5; N-terminal acetyltransferase B complex catalytic subunit NAA20; N-acetyltransferase 5, ARD1 subunit (arrest-defective 1, S. cerevisiae, homolog); NAT5;

Gene ID

51126

mRNA Refseq

NM_016100

Protein Refseq

NP_057184

MIM

610833

UniProt ID

P61599

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Chromosome Location	20p11.23
Function	N-acetyltransferase activity; peptide alpha-N-acetyltransferase activity; transferase activity, transferring acyl groups;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA