

Recombinant Human VEGFA protein (HPLC-verified)

Cat. No. VEGFA-86H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human VEGFA protein (Ala27-Arg191)(HPLC-verified) carries no tag and was expressed in human 293 cells (HEK293).
Species	Human
Source	HEK293
ProteinLength	165
Description	VEGF165 is the most abundant splice variant of VEGF-A. VEGF165 is produced by a number of cells including endothelial cells, macrophages and T cells. VEGF165 is involved in angiogenesis, vascular endothelial cell survival, growth, migration and vascular permeability. VEGF gene expression is induced by hypoxia, inflammatory cytokines and oncogenes. VEGF165 binds to heparan sulfate and is retained on the cell surface and in the extracellular matrix. VEGF165 binds to the receptor tyrosine kinases, VEGFR1 and VEGFR2. VEGF165 is the only splice variant that binds to co-receptors NRP-1 and NRP-2 that function to enhance VEGFR2 signaling. Binding of VEGF165 to VEGFR1 and VEGFR2 leads to activation of the PI3K/AKT, p38 MAPK, FAK and paxillin. VEGF plays a key role in tumor angiogenesis in many cancers.
Form	Lyophilized from 0.22 um filtered solution in PBS, pH7.4, 10% trehalose.
Molecular Mass	The protein has a calculated MW of 19.0 kDa (monomer). As a result of glycosylation, the protein migrates as 24 kDa (monomer) under reducing (R) condition, and 43-50 kDa (homodimer) under non-reducing (NR) condition (SDS-PAGE).

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	Less than 1.0 EU per ug by the LAL method.
Purity	>98% as determined by SDS-PAGE.
Storage	For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower. Please avoid repeated freeze-thaw cycles. This product is stable after storage at: -20 centigrade to -70 centigrade for 12 months in lyophilized state; -70 centigrade for 3 months under sterile conditions after reconstitution.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover the entire contents.


GENE INFORMATION

Gene Name	VEGFA
Official Symbol	VEGFA
Synonyms	VEGFA; vascular endothelial growth factor A; vascular endothelial growth factor , VEGF; VEGF A; VPF; vascular permeability factor; VEGF; MVCD1; MGC70609
Gene ID	7422
mRNA Refseq	NM_001025366
Protein Refseq	NP_001020537
MIM	192240


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID
P15692
**Bioactivity-ELISA of
VEGFA-86H**


Immobilized Human VEGF165, Tag Free (HPLC-verified) at 0.01 µg/mL (100 µL/well) can bind Human VEGFR1, His Tag with a linear range of 4-31 ng/mL (QC tested).

**Bioactivity-ELISA of
VEGFA-86H**



Immobilized Human VEGF165, Tag Free (HPLC-verified) at 2 µg/mL (100 µL/well) can bind Biotinylated Human VEGFR2, Avitag,His Tag with a linear range of 10-156 ng/mL (Routinely tested).

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

Bioactivity-ELISA of VEGFA-86H



Immobilized Human VEGF165, Tag Free (HPLC-verified) at 2 µg/mL (100 µL/well) can bind pre-mixed increasing concentrations of Bevacizumab and 0.5 µg/mL (100 µL/well) Biotinylated Human VEGF R2, Avitag, His Tag with a half maximal inhibitory concentration (IC₅₀) of 0.70 µg/mL (Routinely tested).

Bioactivity-SPR of VEGFA-86H



Avastin (Bevacizumab) captured on CM5 chip via ant-human IgG Fc antibodies surface, can bind Human VEGF165, Tag Free (HPLC-verified) with an affinity constant of 0.776 nM as determined in a SPR assay (Routinely tested).


Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

**Bioactivity-Cell
based assay of
VEGFA-86H**


Recombinant Human VEGF165 stimulates proliferation of HUVEC



Human VEGF165, Tag Free (HPLC-verified) stimulates proliferation of human umbilical vein endothelial cells (HUVEC). The ED50 for this effect is 4.216-9.281 ng/ml (Routinely tested).


**Bioactivity-Cell
based assay of
VEGFA-86H**

Inhibition effect of anti-VEGF mAb (Avastin) in VEGF165 proliferation assay




Inhibition assay shows that the proliferation effect of Human VEGF165, Tag Free (HPLC-verified) is inhibited by increasing concentration of anti-VEGF mAb (Avastin). The concentration of VEGF165 used is 20 ng/ml. The ED50 is 0.065-0.229 µg/mL (Routinely tested).

SDS-PAGE of VEGFA-86H



Human VEGF165, Tag Free (HPLC-verified) on SDS-PAGE under reducing (R) and non-reducing (NR) conditions. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 98%.

SEC-HPLC of VEGFA-86H



A SEC-HPLC analysis showing 95% of Human VEGF165, Tag Free (HPLC-verified) present as active homodimers.

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA