

Recombinant Human ICOS protein, hFc-tagged

Cat. No. ICOS-3244H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human ICOS protein(Q9Y6W8-1)(Glu21-Phe141), fused with C-terminal hFc tag, was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	Glu21-Phe141
Tag	C-hFc
Form	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 5% trehalose is added as protectant before lyophilization.
Bio-activity	Immobilized Human B7-H2, His Tag at 1ug/ml (100ul/Well) on the plate. Dose response curve for Human ICOS, hFc Tag with the EC50 of 15.2ng/ml determined by ELISA.
Molecular Mass	The protein has a predicted MW of 40.5 kDa. Due to glycosylation, the protein migrates to 52-60 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE;> 95% as determined by HPLC


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage	Reconstituted protein stable at -80°C for 12 months, 4°C for 1 week. Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Reconstitution	Centrifuge tubes before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
GENE INFORMATION	
Gene Name	ICOS inducible T-cell co-stimulator [Homo sapiens]
Official Symbol	ICOS
Synonyms	ICOS; inducible T-cell co-stimulator; inducible T-cell costimulator; activation inducible lymphocyte immunomediatory molecule; AILIM; CD278; inducible costimulator; activation-inducible lymphocyte immunomediatory molecule; CVID1; MGC39850;
Gene ID	29851
mRNA Refseq	NM_012092
Protein Refseq	NP_036224
MIM	604558
UniProt ID	Q9Y6W8

Tris-Bis PAGE



ELISA Data



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA