

Intestinal Permeability Assay Kit

Cat. No. Kit-2077 **Lot. No.** (See product label)

SPECIFICATION

Product Overview

The intestinal permeability assay kit is based on measuring the ratio of the absorption of two non-metabolized sugars to through the intestines. Test subjects drink a prescribed amount of lactulose and mannitol and the % absorption of these sugars is determined by the amount of excreted lactulose and mannitol measured during the first 6 hours after ingestion. The degree of intestinal permeability is reflected by the ratio of the % absorption of lactulose to % absorption of mannitol. An increase in this ratio indicates increased intestinal permeability since lactulose is only absorbed through intercellular spaces. Lactulose and mannitol are measured in separate assays using the included Lactulose Assay Kit and Mannitol Assay Kit, respectively.

Size

100 tests

Description

INTESTINAL PERMEABILITY is a phenomenon of the gut wall in which leakage of molecules and ions below ~0.4 nm occurs from the gut lumen into blood circulation. This paracellular leakage occurs through tight junctions between epithelial cells. Elevated paracellular leakage has been implicated in many disorders including type 1 and type 2 diabetes, obesity, inflammatory bowel disease, celiac disease, Parkinson's disease and cancer.

Applications

Determination of intestinal permeability (leaky gut syndrome) through measuring lactulose/mannitol ratio.

Target Species

Human

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage	The kit is shipped on ice. Store all components at -20°C upon receiving. Shelf life: 6 months after receipt.
Kit Components	LACTULOSE ASSAY KIT Assay Buffer: 6 mL Standard: 400 µ 15 mM Lactulose Enzyme A: Dried Enzyme B: 120 µ Enzyme Buffer: 150 µ PMS Solution: 1.5 mL MANNITOL ASSAY KIT Assay Buffer: 10 mL Standard: 0.5 mL 20 mM D-Mannitol Enzyme: 120 µ
Detection method	OD565nm
Compatible Sample Types	Urine
Features & Benefits	• Simple and convenient. Both assays require addition of single working reagent and can be completed within 60 minutes. Both assays are performed at room temperature. No 37°C heater is needed. • High-throughput. "Add-mix-read" type assay. Can be readily automated as a high-throughput 96-well plate assay for thousands of samples per day.
Assay time	60 and 30 min
Sensitivity	Detection Limit: D-Mannitol: 7 µM; Lactulose: 3 µM